

UNIVERSAL GRADED SPECHT MODULES FOR CYCLOTOMIC HECKE ALGEBRAS

ALEXANDER S. KLESHCHEV, ANDREW MATHAS, AND ARUN RAM

ABSTRACT. The graded Specht module S^λ for a cyclotomic Hecke algebra comes with a distinguished generating vector $z^\lambda \in S^\lambda$, which can be thought of as a “highest weight vector of weight λ ”. This paper describes the *defining relations* for the Specht module S^λ as a graded module generated by z^λ . The first three relations say precisely what it means for z^λ to be a highest weight vector of weight λ . The remaining relations are homogeneous analogues of the classical *Garnir relations*. The homogeneous Garnir relations, which are *simpler* than the classical ones, are associated with a remarkable family of homogeneous operators on the Specht module which satisfy the braid relations.

1. INTRODUCTION

Let S_d be the symmetric group on d letters. A central role in representation theory of S_d is played by certain $\mathbb{Z}S_d$ -modules S^λ labelled by the partitions λ of d . These modules are called *Specht modules* and their construction goes back to [26, 29, 30]. Specht modules also arise naturally as cell modules in the cellular structure on the group algebra of S_d constructed by Murphy in [23], see [7, 14, 21] for further development of these ideas which will be important in this paper.

It was shown recently by Brundan and the first author [3] that over an arbitrary field F , the group algebra FS_d is explicitly isomorphic to a certain cyclotomic Khovanov-Lauda-Rouquier (KLR) algebra R_d^Λ . The algebra R_d^Λ is $\mathbb{Z}$ -graded, and this grading can be transferred to FS_d using the Brundan-Kleshchev isomorphism. Moreover, in [6], the Specht modules over F were also explicitly graded, which played a crucial role in the graded categorification theorem of [4] generalizing the Ariki’s categorification theorem [1]. We refer the reader to [19] for description of these ideas and further references.

Hu and the second author [11] have completed the picture by constructing a *graded cellular* structure on the group algebra of the symmetric group, so that the graded Specht modules of [6] arise as the corresponding cell modules.

In all constructions above, the Specht module S^λ comes together with a remarkable generating vector $z^\lambda \in S^\lambda$, which can be thought of, informally, as a “highest weight vector of weight λ ”. The goal of this paper is to describe the *defining relations* of the Specht module S^λ over $\mathbb{Z}$ as a

2000 *Mathematics Subject Classification.* 20C08, 20C30, 05E10.

Research supported by the NSF (grant DMS-0654147), the Australian Research Council (DP0986774, DP0986349, DP110100050), an International Visiting Professorship at the University of Sydney, the University of Melbourne, and the Hausdorff Institute for Mathematics. The first author thanks the University of Melbourne and the University of Sydney for hospitality.

graded module over the KLR algebra R_d^Λ generated by z^λ . This idea of presenting Specht modules by generators and relations is responsible for our terminology *universal Specht modules*.

Our homogeneous relations for S^λ are given in Definition 5.9. The first three relations say precisely what we mean by z^λ being a highest weight vector of weight λ . The fourth and final relation is a remarkable family of *homogeneous Garnir relations*, which we consider to be the key innovation of this paper.

We point out that the classical Garnir relations, which go back to [8, 30], are very far from being homogeneous with respect to the gradings under consideration. The classical Garnir relations have the form of an alternating sum of elements of the Specht module corresponding to certain tableaux (being equated to zero).

Even though substantial initial work is required to define the homogeneous Garnir relations, they are actually much *simpler* than the classical ones. For example, if the underlying Lie type of the KLR algebra is A_∞ , which under the isomorphism of [3] corresponds to the case where the field F has characteristic 0, then the homogeneous Garnir relation has the form of just *one* element corresponding to a special Garnir tableaux (being equated to zero). In the case where the Lie type is $A_{p-1}^{(1)}$, which under the isomorphism of [3] corresponds to the field F having characteristic $p > 0$, the homogeneous Garnir relation does look like a sum, but it has roughly p times as few summands as the classical Garnir relation. For the case of the so-called calibrated representations of the affine Hecke algebra in characteristic zero this phenomenon has been known, see for example [24, (5.4)].

Even though so far we have been talking only about the symmetric groups, the story of Specht modules generalizes to all *cyclotomic Hecke algebras*, both degenerate and non-degenerate. This is the generality which we work with throughout this paper.

In section 2 we collect various combinatorial facts and notation. The key notion here is that of the degree of a standard tableau which was first defined in [6]. In section 3, we recall the definition of the affine and cyclotomic KLR algebras and define “permutation modules” for these algebras using induction from one-dimensional modules of the parabolic subalgebras in the affine setting.

In the crucial section 4, we define certain elements which we call *block intertwiners*. These intertwiners will later be fed into the definition of the homogeneous Garnir relations. They permute blocks (or bricks) of size e , where e can be thought of as the analogue of the characteristic of the ground field when working with Specht modules for the symmetric groups, and this part of the story is trivial when $e = 0$. The block intertwiners τ_r are defined in terms of products of the large number of the KLR generators. The KLR generators do not satisfy Coxeter relations, so we find it truly remarkable that the brick intertwiners τ_r do! See the key Theorem 4.12.

In section 5, we define (row) Garnir relations and *universal (row) Specht modules* S^λ for the algebra R_d^Λ by generators and relations, see Definition 5.9. Our next goal is to prove that if we identify the cyclotomic KLR algebra R_d^Λ with the cyclotomic Hecke algebra H_d^Λ via the Brundan-Kleshchev isomorphism, which is only valid over a field, then the universal Specht modules

are identified with the usual graded Specht modules of [6]. This is done in section 6.

Section 7 develops the parallel story for the column Specht modules S_λ , which turn out to be dual to the row Specht modules S^λ . Accidentally, what we call a column Specht module S_λ is what was called a Specht module in James' book [13].

The final section 8 contains two applications. One is the description of Specht modules for higher level cyclotomic Hecke algebras as modules induced from Specht modules of level 1, see Theorem 8.2. In fact, these induced modules were sometimes taken as a definition of Specht modules for higher levels, which was problematic because the connection with the Specht modules as cell modules had not been established in full generality before.

Our second application is a generalization of the famous very useful result from James' book [13, Theorem 8.15] for symmetric groups:

$$(S^\lambda)^* \cong S^{\lambda'} \otimes \text{sgn}.$$

The analogue of this is proved for arbitrary cyclotomic Hecke algebras in the graded setting, see Theorem 8.5.

2. COMBINATORICS

2.1. Lie theoretic notation. Let $e \in \{0, 2, 3, 4, \dots\}$ and $I := \mathbb{Z}/e\mathbb{Z}$. Let Γ be the quiver with vertex set I , and a directed edge from i to j if $j = i - 1$ (the orientation differs from the one in [3, 11]). Thus Γ is a quiver of type A_∞ if $e = 0$ or $A_{e-1}^{(1)}$ if $e > 0$. The corresponding *Cartan matrix* $(a_{i,j})_{i,j \in I}$ is defined by

$$a_{i,j} := \begin{cases} 2 & \text{if } i = j, \\ 0 & \text{if } j \neq i, i \pm 1, \\ -1 & \text{if } i \rightarrow j \text{ or } i \leftarrow j, \\ -2 & \text{if } i \rightleftarrows j. \end{cases} \quad (2.1)$$

(The case $a_{i,j} = -2$ only occurs if $e = 2$.)

Following [15], let $(\mathfrak{h}, \Pi, \Pi^\vee)$ be a realization of the Cartan matrix $(a_{ij})_{i,j \in I}$, so we have the simple roots $\{\alpha_i \mid i \in I\}$, the fundamental dominant weights $\{\Lambda_i \mid i \in I\}$, and the normalized invariant form $(\cdot, \cdot)$ such that

$$(\alpha_i, \alpha_j) = a_{ij}, \quad (\Lambda_i, \alpha_j) = \delta_{ij} \quad (i, j \in I).$$

If $e > 0$, the *null-root* is

$$\delta := \alpha_0 + \alpha_1 + \dots + \alpha_{e-1}. \quad (2.2)$$

Let P_+ be the set of dominant integral weights, and $Q_+ := \bigoplus_{i \in I} \mathbb{Z}_{\geq 0} \alpha_i$ the positive part of the root lattice. For $\alpha \in Q_+$ let $\text{ht}(\alpha)$ be the *height* of α . That is, $\text{ht}(\alpha)$ is the sum of the coefficients when α is expanded in terms of the α_i 's.

Let $\mathfrak{S}_d$ be the *symmetric group* on d letters and let $s_r = (r, r+1)$, for $1 \leq r < d$, be the simple transpositions of $\mathfrak{S}_d$. Then $\mathfrak{S}_d$ acts from the left on the set I^d by place permutations. If $\mathbf{i} = (i_1, \dots, i_d) \in I^d$ then its *weight* is $|\mathbf{i}| := \alpha_{i_1} + \dots + \alpha_{i_d} \in Q_+$. Then the $\mathfrak{S}_d$ -orbits on I^d are the sets

$$I^\alpha := \{\mathbf{i} \in I^d \mid \alpha = |\mathbf{i}|\}$$

parametrized by all $\alpha \in Q_+$ of height d .

Throughout the paper, we fix a positive integer l , referred to as the *level*, and an ordered l -tuple

$$\kappa = (k_1, \dots, k_l) \in I^l. \quad (2.3)$$

Define the dominant weight Λ (of level l) as follows:

$$\Lambda = \Lambda(\kappa) := \Lambda_{k_1} + \dots + \Lambda_{k_l} \in P_+. \quad (2.4)$$

Finally, for $\alpha \in Q_+$, define the *defect* of α (relative to Λ) to be


$$\text{def}(\alpha) = (\Lambda, \alpha) - \frac{1}{2}(\alpha, \alpha). \quad (2.5)$$

2.2. Partitions. Recall that in (2.3) we have fixed a level l and an l -tuple $\kappa = (k_1, \dots, k_l)$. An l -*multipartition* of d is an ordered l -tuple of partitions $\mu = (\mu^{(1)}, \dots, \mu^{(l)})$ such that $\sum_{m=1}^l |\mu^{(m)}| = d$. We call $\mu^{(m)}$ the m th *component* of μ . Let $\mathcal{P}_d^\kappa$ be the set of all l -multipartitions of d and put $\mathcal{P}^\kappa := \bigsqcup_{d \geq 0} \mathcal{P}_d^\kappa$. Of course, $\mathcal{P}^\kappa$ only depends on l , and not on κ , but as soon as we consider residues of nodes of multipartitions, the dependence on κ becomes crucial.

The *Young diagram* of the multipartition $\mu = (\mu^{(1)}, \dots, \mu^{(l)}) \in \mathcal{P}^\kappa$ is

$$\{(a, b, m) \in \mathbb{Z}_{>0} \times \mathbb{Z}_{>0} \times \{1, \dots, l\} \mid 1 \leq b \leq \mu_a^{(m)}\}.$$

The elements of this set are the *nodes* of μ . More generally, a *node* is any element of $\mathbb{Z}_{>0} \times \mathbb{Z}_{>0} \times \{1, \dots, l\}$. Usually, we identify the multipartition μ with its Young diagram and visualize it as a column vector of Young diagrams. For example, $((3, 1), \emptyset, (4, 2))$ is the Young diagram



To each node $A = (a, b, m)$ we associate its *residue*, which is the following element of $I = \mathbb{Z}/e\mathbb{Z}$:

$$\text{res } A = \text{res}^\kappa A = k_m + (b - a) \pmod{e}. \quad (2.6)$$

An i -*node* is a node of residue i . Define the *residue content* of μ to be

$$\text{cont}(\mu) := \sum_{A \in \mu} \alpha_{\text{res } A} \in Q_+. \quad (2.7)$$

Denote

$$\mathcal{P}_\alpha^\kappa := \{\mu \in \mathcal{P}^\kappa \mid \text{cont}(\mu) = \alpha\} \quad (\alpha \in Q_+).$$

A node $A \in \mu$ is a *removable node* (of μ) if $\mu \setminus \{A\}$ is (the diagram of) a multipartition. A node $B \notin \mu$ is an *addable node* (for μ) if $\mu \cup \{B\}$ is a multipartition. We use the notation

$$\mu_A := \mu \setminus \{A\}, \quad \mu^B := \mu \cup \{B\}.$$

Let $\mu, \nu \in \mathcal{P}_d^\kappa$. Then μ *dominates* ν , and we write $\mu \trianglerighteq \nu$, if

$$\sum_{a=1}^{m-1} |\mu^{(a)}| + \sum_{b=1}^c \mu_b^{(m)} \geq \sum_{a=1}^{m-1} |\nu^{(a)}| + \sum_{b=1}^c \nu_b^{(m)}$$

for all $1 \leq m \leq l$ and $c \geq 1$. In other words, μ is obtained from ν by moving nodes up in the diagram.

We define

$$\kappa' := (-k_l, \dots, -k_1). \quad (2.8)$$

Now, let $\mu = (\mu^{(1)}, \dots, \mu^{(l)}) \in \mathcal{P}^\kappa$. The *conjugate* of μ is the multipartition

$$\mu' = (\mu^{(l)'}, \dots, \mu^{(1)'}) \in \mathcal{P}^{\kappa'},$$

where each $\mu^{(m)'}$ is the partition conjugate to $\mu^{(m)}$ in the usual sense, that is, $\mu^{(m)'}$ is obtained by swapping the rows and columns of $\mu^{(m)}$.

2.3. Tableaux. Let $\mu = (\mu^{(1)}, \dots, \mu^{(l)}) \in \mathcal{P}_d^\kappa$. A μ -*tableau* $T = (T^{(1)}, \dots, T^{(l)})$ is obtained from the diagram of μ by inserting the integers $1, \dots, d$ into the nodes, allowing no repeats. For each $m = 1, \dots, l$, $T^{(m)}$ is a $\mu^{(m)}$ -tableau, called the m th *component* of T . If the node $A = (a, b, m) \in \mu$ is occupied by the integer r in T then we write $r = T(a, b, m)$ and set $\text{res}_T(r) = \text{res } A$. The *residue sequence* of T is

$$\mathbf{i}(T) = \mathbf{i}^\kappa(T) = (i_1, \dots, i_d) \in I^d, \quad (2.9)$$

where $i_r = \text{res}_T(r)$ is the residue of the node occupied by r in T ($1 \leq r \leq d$).

A μ -tableau T is *row-strict* (resp. *column-strict*) if its entries increase from left to right (resp. from top to bottom) along the rows (resp. columns) of each component of T . A μ -tableau T is *standard* if it is row- and column-strict. Let $\text{St}(\mu)$ be the set of standard μ -tableaux.

Let T be a μ -tableau and suppose that $1 \leq r \neq s \leq d$ and that $r = T(a_1, b_1, m_1)$ and that $s = T(a_2, b_2, m_2)$. We write $r \nearrow_T s$ if $m_1 = m_2$, $a_1 > a_2$, and $b_1 < b_2$; informally, r and s are in the same component and s is strictly to the north-east of r within that component. The symbols $\rightarrow_T$, $\searrow_T$, $\downarrow_T$ have the similar obvious meanings. For example, $r \downarrow_T s$ means that r and s are located in the same column of the same component of T and that s is in a strictly lower row of T than r .

Let $\mu \in \mathcal{P}^\kappa$, $i \in I$, A be a removable i -node and B be an addable i -node of μ . We set

$$d_A(\mu) = \# \left\{ \begin{array}{c} \text{addable } i\text{-nodes of } \mu \\ \text{strictly below } A \end{array} \right\} - \# \left\{ \begin{array}{c} \text{removable } i\text{-nodes of } \mu \\ \text{strictly below } A \end{array} \right\}, \quad (2.10)$$

and

$$d^B(\mu) = \# \left\{ \begin{array}{c} \text{addable } i\text{-nodes of } \mu \\ \text{strictly above } B \end{array} \right\} - \# \left\{ \begin{array}{c} \text{removable } i\text{-nodes of } \mu \\ \text{strictly above } B \end{array} \right\}. \quad (2.11)$$

Given $\mu \in \mathcal{P}_d^\kappa$ and $T \in \text{St}(\mu)$, the *degree* of T is defined in [6, section 3.5] inductively as follows. If $d = 0$, then T is the empty tableau $\emptyset$, and we set $\deg(T) := 0$. Otherwise, let A be the node occupied by d in T . Let $T_{<d} \in \text{St}(\mu_A)$ be the tableau obtained by removing this node and set

$$\deg(T) := d_A(\mu) + \deg(T_{<d}). \quad (2.12)$$

Similarly, define a dual notion of *codegree* by

$$\text{codeg}(\emptyset) := 0, \quad \text{codeg}(T) := d^A(\mu_A) + \text{codeg}(T_{<d}). \quad (2.13)$$

The definition of the degree and codegree of a tableau depend on the residues and so, ultimately, they depend on κ by (2.6). We write $\deg^\kappa(T)$ and $\text{codeg}^\kappa(T)$ when we wish to emphasize this dependence.

By [6, Lemma 3.12], using codegree instead of degree for a tableau leads only to a negation and “global shift” by the defect: more precisely, we have

$$\deg(T) + \text{codeg}(T) = \text{def}(\alpha) \quad (T \in \text{St}(\mu), \mu \in \mathcal{P}_\alpha^\kappa). \quad (2.14)$$

The group $\mathfrak{S}_d$ acts on the set of μ -tableaux from the left by acting on the entries of the tableaux. Let T^μ be the μ -tableau in which the numbers $1, 2, \dots, d$ appear in order from left to right along the successive rows, working from top row to bottom row. Let T_μ be the μ -tableau in which the numbers $1, 2, \dots, d$ appear in from top to bottom along the successive columns, working from the leftmost column to the rightmost column within a component and moving from the l th component up to the first component.

For example, if $\mu = ((3, 1), (2, 2))$ then

$$T^\mu = \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 4 & & \\ \hline \end{array} \quad \text{and} \quad T_\mu = \begin{array}{|c|c|c|} \hline 5 & 7 & 8 \\ \hline 6 & & \\ \hline \end{array}.$$

$$\begin{array}{|c|c|} \hline 5 & 6 \\ \hline 7 & 8 \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & 4 \\ \hline \end{array}.$$

Set

$$\mathbf{i}^\mu := \mathbf{i}(T^\mu) \quad \text{and} \quad \mathbf{i}_\mu := \mathbf{i}(T_\mu). \quad (2.15)$$

For each μ -tableau T define permutations w^T and w_T in $\mathfrak{S}_d$ by the equations

$$w^T T^\mu = T = w_T T_\mu. \quad (2.16)$$

If $T = (T^{(1)}, \dots, T^{(l)}) \in \text{St}(\mu)$ then the *conjugate* of T is the standard μ' -tableau $T' = (T^{(l)'}, \dots, T^{(1)'})$, where $T^{(m)'} is the $\mu^{(m)'}-tableau obtained by swapping the rows and columns of $T^{(m)}$, for $1 \leq m \leq l$. For example, $(T^\mu)' = T_{\mu'}$.$$

2.4. Bruhat order. Let ℓ be the length function on $\mathfrak{S}_d$ with respect to the Coxeter generators $s_1, s_2, \dots, s_{d-1}$. Let $\leq$ be the *Bruhat order* on $\mathfrak{S}_d$ (so that $1 \leq w$ for all $w \in \mathfrak{S}_d$). Define a related partial order on $\text{St}(\mu)$ as follows: if $S, T \in \text{St}(\mu)$ then

$$S \trianglelefteq T \quad \text{if and only if} \quad w_S \leq w_T. \quad (2.17)$$

If $S \trianglelefteq T$ then we also write $T \trianglerighteq S$. If $S \trianglelefteq T$ and $S \neq T$ we write $S \triangleleft T$ and $T \triangleright S$.

Observe that if $T \in \text{St}(\mu)$ then $T_\mu \trianglelefteq T \trianglelefteq T^\mu$. There is a similar connection between the relation $w^S \leq w^T$ and the corresponding tableaux. To describe this, recall conjugate multipartitions and tableaux.

Lemma 2.18. *Suppose that $\mu \in \mathcal{P}_d^\kappa$ and that $S, T \in \text{St}(\mu)$. Then:*

- (i) $w_T = w^{T'}$;
- (ii) $T \trianglelefteq S$ if and only if $w^T \geq w^S$;
- (iii) $w_{T^\mu} = (w^T)^{-1} w_T$ and $w^{T^\mu} = w_T^{-1} w^T$ with $\ell(w_{T^\mu}) = \ell(w^T) + \ell(w_T)$ and $\ell(w^{T^\mu}) = \ell(w_T) + \ell(w^T)$.

Proof. (i) Observe that $(T^\mu)' = T_{\mu'}$, $(T_\mu)' = T^{\mu'}$ and $\text{St}(\mu) = \{T \mid T' \in \text{St}(\mu')\}$. Now, conjugating the equation $T = w^T T^\mu$ shows that $w_{T'} = w^T$, for $T \in \text{St}(\mu)$.

(ii) If $U \in \text{St}(\mu)$ and $1 \leq k \leq d$, let $U_{\leq k}$ be the subtableau of T containing the entries $1, 2, \dots, k$. Then it follows from [21, Theorem 3.8] that $S \trianglerighteq T$ if

and only if the shape of $\mathbf{S}_{\leq k}$ dominates the shape of $\mathbf{T}_{\leq k}$ for all $1 \leq k \leq d$. So $\mathbf{T} \trianglelefteq \mathbf{S}$ if and only if $\mathbf{S}' \trianglelefteq \mathbf{T}'$. Therefore, by part (i) and (2.17), we get

$$\mathbf{T} \trianglelefteq \mathbf{S} \iff \mathbf{S}' \trianglelefteq \mathbf{T}' \iff w_{\mathbf{S}'} \leq w_{\mathbf{T}'} \iff w^{\mathbf{S}} \leq w^{\mathbf{T}}.$$

(iii) Since $w_{\mathbf{T}}\mathbf{T}_{\mu} = \mathbf{T} = w^{\mathbf{T}}\mathbf{T}^{\mu}$, we have $w_{\mathbf{T}^{\mu}}\mathbf{T}_{\mu} = (w^{\mathbf{T}})^{-1}w_{\mathbf{T}}\mathbf{T}_{\mu}$ which implies that $w_{\mathbf{T}^{\mu}} = (w^{\mathbf{T}})^{-1}w_{\mathbf{T}}$. Since $\mathbf{T}_{\mu} \trianglelefteq \mathbf{T} \trianglelefteq \mathbf{T}^{\mu}$ we obtain $\ell(w_{\mathbf{T}^{\mu}}) = \ell(w^{\mathbf{T}}) + \ell(w_{\mathbf{T}})$ using the description of the Bruhat order given in (ii). The remaining claims are proved similarly. $\square$

We will also need the following result.

Lemma 2.19. [6, Lemma 3.7] *Suppose that $\mu \in \mathcal{P}_d^{\kappa}$, $\mathbf{T} \in \text{St}(\mu)$, and $1 \leq r < d$ such that $r \downarrow_{\mathbf{T}} r+1$ or $r \rightarrow_{\mathbf{T}} r+1$. Suppose that $\mathbf{S} \in \text{St}(\mu)$ and $\mathbf{S} \triangleright_{s_r} \mathbf{T}$. Then $\mathbf{S} \triangleright \mathbf{T}$.*

3. KLR ALGEBRAS AND PERMUTATION MODULES

Throughout this paper a *graded algebra* will mean a $\mathbb{Z}$ -graded algebra and a *graded module* will be a $\mathbb{Z}$ -graded module. If A is a graded algebra then $A\text{-Mod}$ is the category of finitely generated graded (left) A -modules with degree preserving maps. We use the standard notation of graded representation theory. In particular, if $M = \bigoplus_{d \in \mathbb{Z}} M_d$ then $v \in M_d$ is *homogeneous of degree $d = \deg v$* . Further, if $n \in \mathbb{Z}$ then $M\langle n \rangle$ is the graded module obtained by shifting the grading on M up by n so that $M\langle n \rangle_d = M_{d-n}$.

3.1. KLR algebras. Let $\mathcal{O}$ be a commutative ring with identity and $\alpha \in Q_+$. Recall from [16, 17, 25] that the (affine) *Khovanov-Lauda-Rouquier algebra*, or KLR algebra, $R_{\alpha} = R_{\alpha}(\mathcal{O})$, is defined to be the unital $\mathcal{O}$ -algebra generated by the elements

$$\{e(\mathbf{i}) \mid \mathbf{i} \in I^{\alpha}\} \cup \{y_1, \dots, y_d\} \cup \{\psi_1, \dots, \psi_{d-1}\}, \quad (3.1)$$

subject only to the following relations:

$$e(\mathbf{i})e(\mathbf{j}) = \delta_{\mathbf{i}, \mathbf{j}}e(\mathbf{i}); \quad \sum_{\mathbf{i} \in I^{\alpha}} e(\mathbf{i}) = 1; \quad (3.2)$$

$$y_r e(\mathbf{i}) = e(\mathbf{i}) y_r; \quad \psi_r e(\mathbf{i}) = e(s_r \mathbf{i}) \psi_r; \quad (3.3)$$

$$y_r y_s = y_s y_r; \quad (3.4)$$

$$\psi_r y_s = y_s \psi_r \quad \text{if } s \neq r, r+1; \quad (3.5)$$

$$\psi_r \psi_s = \psi_s \psi_r \quad \text{if } |r-s| > 1; \quad (3.6)$$

$$\psi_r y_{r+1} e(\mathbf{i}) = (y_r \psi_r + \delta_{i_r, i_{r+1}}) e(\mathbf{i}) \quad (3.7)$$

$$y_{r+1} \psi_r e(\mathbf{i}) = (\psi_r y_r + \delta_{i_r, i_{r+1}}) e(\mathbf{i}) \quad (3.8)$$

$$\psi_r^2 e(\mathbf{i}) = \begin{cases} 0 & \text{if } i_r = i_{r+1}, \\ e(\mathbf{i}) & \text{if } i_{r+1} \neq i_r, i_r \pm 1, \\ (y_{r+1} - y_r) e(\mathbf{i}) & \text{if } i_r \rightarrow i_{r+1}, \\ (y_r - y_{r+1}) e(\mathbf{i}) & \text{if } i_r \leftarrow i_{r+1}, \\ (y_{r+1} - y_r)(y_r - y_{r+1}) e(\mathbf{i}) & \text{if } i_r \rightleftharpoons i_{r+1}; \end{cases} \quad (3.9)$$

$$\psi_r \psi_{r+1} \psi_r e(\mathbf{i}) = \begin{cases} (\psi_{r+1} \psi_r \psi_{r+1} + 1) e(\mathbf{i}) & \text{if } i_{r+2} = i_r \rightarrow i_{r+1}, \\ (\psi_{r+1} \psi_r \psi_{r+1} - 1) e(\mathbf{i}) & \text{if } i_{r+2} = i_r \leftarrow i_{r+1}, \\ (\psi_{r+1} \psi_r \psi_{r+1} - 2y_{r+1} + y_r + y_{r+2}) e(\mathbf{i}) & \text{if } i_{r+2} = i_r \rightleftharpoons i_{r+1}, \\ \psi_{r+1} \psi_r \psi_{r+1} e(\mathbf{i}) & \text{otherwise.} \end{cases} \quad (3.10)$$

Recall from (2.4) that we have fixed $\Lambda = \Lambda(\kappa) \in P_+$. The corresponding *cyclotomic KLR algebra* $R_\alpha^\Lambda = R_\alpha^\Lambda(\mathcal{O})$ is generated by the same elements (3.1) subject only to the relations (3.2)–(3.10) and the additional *cyclotomic relations*

$$y_1^{(\Lambda, \alpha_{i_1})} e(\mathbf{i}) = 0 \quad (\mathbf{i} = (i_1, \dots, i_d) \in I^\alpha). \quad (3.11)$$

Thus R_α^Λ is the quotient of R_α by the relations (3.11).

The algebras R_α and R_α^Λ have $\mathbb{Z}$ -gradings determined by setting $e(\mathbf{i})$ to be of degree 0, y_r of degree 2, and $\psi_r e(\mathbf{i})$ of degree $-a_{i_r, i_{r+1}}$ for all r and $\mathbf{i} \in I^\alpha$.

Note that $R_\alpha(\mathbb{Z}) \otimes_{\mathbb{Z}} \mathcal{O} \cong R_\alpha(\mathcal{O})$ and $R_\alpha^\Lambda(\mathbb{Z}) \otimes_{\mathbb{Z}} \mathcal{O} \cong R_\alpha^\Lambda(\mathcal{O})$. In this paper $\mathcal{O}$ will usually be $\mathbb{Z}$ or a field F .

3.2. Graded duality. Let $\alpha \in Q_+$ be of height d . It is easy to check using generators and relations that there exists a homogeneous algebra anti-involution

$$\tau : R_\alpha \longrightarrow R_\alpha, \quad e(\mathbf{i}) \mapsto e(\mathbf{i}), \quad y_r \mapsto y_r, \quad \psi_s \mapsto \psi_s. \quad (3.12)$$

for all $\mathbf{i} \in I^\alpha$, $1 \leq r \leq d$, and $1 \leq s < d$. Note that τ factors through to an anti-involution of the cyclotomic quotient R_α^Λ , which we also denote by τ .

If $M = \bigoplus_{d \in \mathbb{Z}} M_d$ is a finite rank graded R_α -module, then the *graded dual* M^* is the graded $\mathcal{O}$ -module such that $(M^*)_d := \text{Hom}_{\mathcal{O}}(M_{-d}, \mathcal{O})$, for all $d \in \mathbb{Z}$, and where the R_α -action is given by $(xf)(m) = f(\tau(x)m)$, for all $f \in M^*, m \in M, x \in R_\alpha$.

3.3. The sign map. For $\mathbf{i} = (i_1, \dots, i_d) \in I^d$, set

$$-\mathbf{i} := (-i_1, \dots, -i_d). \quad (3.13)$$

If $\alpha = \sum_{i \in I} a_i \alpha_i \in Q_+$, then define

$$\alpha' = \sum_{i \in I} a_i \alpha_{-i}.$$

We clearly have $\alpha' \in Q_+$ and $\text{ht}(\alpha') = \text{ht}(\alpha)$. Moreover, $\mathbf{i} \in I^\alpha$ if and only if $-\mathbf{i} \in I^{\alpha'}$. Now, inspecting the relations, there is a unique homogeneous algebra isomorphism

$$\mathbf{sgn} : R_\alpha \longrightarrow R_{\alpha'}, \quad e(\mathbf{i}) \mapsto e(-\mathbf{i}), \quad y_r \mapsto -y_r, \quad \psi_s \mapsto -\psi_s \quad (3.14)$$

for all $\mathbf{i} \in I^\alpha$, $1 \leq r \leq d$, and $1 \leq s < d$, where $d = \text{ht}(\alpha)$.

Recall $\kappa = (\kappa_1, \dots, \kappa_l)$ from (2.3) and $\kappa' = (-\kappa_l, \dots, -\kappa_1)$ from (2.8). Then, as in (2.4), κ' determines the dominant weight

$$\Lambda' = \Lambda(\kappa') = \Lambda_{-\kappa_l} + \dots + \Lambda_{-\kappa_1} \in P_+.$$

Equivalently, if $\Lambda = \sum_{i \in I} l_i \Lambda_i$, then $\Lambda' = \sum_{i \in I} l_i \Lambda_{-i}$.

The algebra R_α^Λ is the quotient of R_α by the cyclotomic relations (3.11). Applying the involution $\mathbf{sgn}$ to (3.11) we obtain

$$0 = \mathbf{sgn}\left(y_1^{(\Lambda, \alpha_{i_1})} e(\mathbf{i})\right) = \pm y_1^{(\Lambda, \alpha_{i_1})} e(-\mathbf{i}) = \pm y_1^{(\Lambda', \alpha_{-i_1})} e(-\mathbf{i}),$$

where the right hand side is, up to sign, the cyclotomic relation for $R_{\alpha'}^{\Lambda'}$. Hence $\mathbf{sgn}$ factors through to a graded algebra isomorphism

$$\mathbf{sgn} : R_\alpha^\Lambda \xrightarrow{\sim} R_{\alpha'}^{\Lambda'}.$$

The isomorphism $\mathbf{sgn}$ induces equivalences

$$R_{\alpha'}\text{-Mod} \xrightarrow{\sim} R_{\alpha}\text{-Mod} \quad \text{and} \quad R_{\alpha'}^{\Lambda'}\text{-Mod} \xrightarrow{\sim} R_{\alpha}^{\Lambda}\text{-Mod}$$

of the corresponding categories of graded modules. These equivalences send the $R_{\alpha'}$ -module M to the R_{α} -module $M^{\mathbf{sgn}}$, where $M^{\mathbf{sgn}} = M$ as a graded vector space and where the R_{α} -action on $M^{\mathbf{sgn}}$ is given by $a \cdot m = \mathbf{sgn}(a)m$, for $a \in R_{\alpha}$ and $m \in M^{\mathbf{sgn}}$.

3.4. Basis Theorem. Suppose that $\alpha \in Q_+$ is of height d . For the rest of this paper we fix a *preferred reduced decomposition* $w = s_{r_1} \dots s_{r_m}$ for each element $w \in \mathfrak{S}_d$, where $m \geq 0$ is as small as possible and $1 \leq r_1, \dots, r_m < d$. Define the elements

$$\psi_w := \psi_{r_1} \dots \psi_{r_m} \in R_{\alpha} \quad (w \in \mathfrak{S}_d).$$

In general, ψ_w depends on the choice of a preferred reduced decomposition of w , but:

Proposition 3.15. *Suppose that $\mathbf{i} \in I^{\alpha}$, and*

$$w = s_{t_1} \dots s_{t_m} = s_{r_1} \dots s_{r_m}$$

are two reduced decompositions of an element $w \in \mathfrak{S}_d$. Then in R_{α} , we have

$$\psi_{t_1} \dots \psi_{t_m} e(\mathbf{i}) = \psi_{r_1} \dots \psi_{r_m} e(\mathbf{i}) + X,$$

where X is a linear combination of elements of the form $\psi_u f(y) e(\mathbf{i})$ such that $u < w$, $f(y)$ is a polynomial in $y_1, \dots, y_d$, and

$$\deg(\psi_u f(y) e(\mathbf{i})) = \deg(\psi_{r_1} \dots \psi_{r_m} e(\mathbf{i})) = \deg(\psi_{t_1} \dots \psi_{t_m} e(\mathbf{i})).$$

Proof. This is proved in [6, Proposition 2.5] for corresponding elements of the cyclotomic KLR algebra R_{α}^{Λ} . As the argument in [6] does not use the relation (3.11) the result holds in R_{α} . $\square$

Suppose now that $\mu \in \mathcal{P}_{\alpha}^{\kappa}$ and that $\mathbf{T}$ is a μ -tableau. In (2.16) we defined the permutations $w_{\mathbf{T}}, w^{\mathbf{T}} \in \mathfrak{S}_d$. Define

$$\psi^{\mathbf{T}} := \psi_{w^{\mathbf{T}}} \quad \text{and} \quad \psi_{\mathbf{T}} := \psi_{w_{\mathbf{T}}}. \quad (3.16)$$

These elements will be used to produce bases of various modules below.

By (3.6), there is one important case where the elements ψ_w are independent of the choice of preferred reduced decomposition of w . An element $w \in \mathfrak{S}_d$ is *fully commutative* if one can go from any reduced decomposition of w to any other reduced decomposition of w using only the commuting braid relations; that is, the relations of the form $s_r s_t = s_t s_r$, for $|r - t| > 1$. We refer the reader to [27] for more details on fully commutative elements. We record the following easy result for future reference:

Lemma 3.17. *Suppose that $1 \leq s < k$ and let $\mathcal{D}$ be the set of the minimal length left coset representatives of the parabolic subgroup $\mathfrak{S}_s \times \mathfrak{S}_{k-s}$ in the symmetric group $\mathfrak{S}_k$. Then every element of $\mathcal{D}$ is fully commutative.*

In general we have the following important result:

Theorem 3.18. [16, Theorem 2.5], [25, Theorem 3.7] *Let $\alpha \in Q_+$. Then*

$$\{\psi_w y_1^{m_1} \dots y_d^{m_d} e(\mathbf{i}) \mid w \in \mathfrak{S}_d, m_1, \dots, m_d \in \mathbb{Z}_{\geq 0}, \mathbf{i} \in I^{\alpha}\}$$

is an $\mathcal{O}$ -basis of R_{α} .

3.5. Induction and restriction for affine KLR algebras. Given $\alpha, \beta \in Q_+$, we set $R_{\alpha, \beta} := R_\alpha \otimes R_\beta$, viewed as an algebra in the usual way. Let $M \boxtimes N$ be the outer tensor product of the R_α -module M and the R_β -module N . There is an obvious injective homogeneous (non-unital) algebra homomorphism $R_{\alpha, \beta} \hookrightarrow R_{\alpha+\beta}$ mapping $e(\mathbf{i}) \otimes e(\mathbf{j})$ to $e(\mathbf{ij})$, where $\mathbf{ij}$ is the concatenation of the two sequences. The image of the identity element of $R_{\alpha, \beta}$ under this map is $e_{\alpha, \beta} := \sum_{\mathbf{i} \in I^\alpha, \mathbf{j} \in I^\beta} e(\mathbf{ij})$. Let $\text{Ind}_{\alpha, \beta}^{\alpha+\beta}$ and $\text{Res}_{\alpha, \beta}^{\alpha+\beta}$ be the corresponding induction and restriction functors between the corresponding categories of *graded* modules:

$$\begin{aligned} \text{Ind}_{\alpha, \beta}^{\alpha+\beta} &:= R_{\alpha+\beta} e_{\alpha, \beta} \otimes_{R_{\alpha, \beta}} ? : R_{\alpha, \beta}\text{-Mod} \rightarrow R_{\alpha+\beta}\text{-Mod}, \\ \text{Res}_{\alpha, \beta}^{\alpha+\beta} &:= e_{\alpha, \beta} R_{\alpha+\beta} \otimes_{R_{\alpha+\beta}} ? : R_{\alpha+\beta}\text{-Mod} \rightarrow R_{\alpha, \beta}\text{-Mod}. \end{aligned}$$

These have obvious generalizations to $n \geq 2$ factors:

$$\begin{aligned} \text{Ind}_{\beta_1, \dots, \beta_n}^{\beta_1 + \dots + \beta_n} &: R_{\beta_1, \dots, \beta_n}\text{-Mod} \rightarrow R_{\beta_1 + \dots + \beta_n}\text{-Mod}, \\ \text{Res}_{\beta_1, \dots, \beta_n}^{\beta_1 + \dots + \beta_n} &: R_{\beta_1 + \dots + \beta_n}\text{-Mod} \rightarrow R_{\beta_1, \dots, \beta_n}\text{-Mod}. \end{aligned}$$

The functor $\text{Res}_{\beta_1, \dots, \beta_n}^{\beta_1 + \dots + \beta_n}$ is left multiplication by the idempotent $e_{\beta_1, \dots, \beta_n}$, so it is exact and sends finite dimensional modules to finite dimensional modules. The functor $\text{Ind}_{\beta_1, \dots, \beta_n}^{\beta_1 + \dots + \beta_n}$ is left adjoint to $\text{Res}_{\beta_1, \dots, \beta_n}^{\beta_1 + \dots + \beta_n}$. Moreover, $R_{\beta_1 + \dots + \beta_n} e_{\beta_1, \dots, \beta_n}$ is a free graded right $R_{\beta_1, \dots, \beta_n}$ -module of finite rank, so $\text{Ind}_{\beta_1, \dots, \beta_n}^{\beta_1 + \dots + \beta_n}$ sends finite dimensional graded modules to finite dimensional graded modules. Finally, if $M_a \in R_{\beta_a}\text{-Mod}$, for $a = 1, \dots, n$, we define

$$M_1 \circ \dots \circ M_n := \text{Ind}_{\beta_1, \dots, \beta_n}^{\beta_1 + \dots + \beta_n} M_1 \boxtimes \dots \boxtimes M_n. \quad (3.19)$$

3.6. Permutation Modules $M(\vec{s})$. For $i \in I$, and $N \in \mathbb{Z}_{\geq 1}$, let $\mathbf{s}(i, N) \in I^N$ be the tuple $(j_1, \dots, j_N)$ with $j_r = i + r - 1 \pmod{e}$. In other words, $\mathbf{s}(i, N)$ is the *segment* of length N starting at i . Similarly, if $N \in \mathbb{Z}_{< 0}$ define $\mathbf{s}(i, N) \in I^{-N}$ be the tuple $(j_1, \dots, j_{-N})$ with $j_r = i - r + 1 \pmod{e}$. For example $\mathbf{s}(0, e) = (0, 1, \dots, e-1)$ and $\mathbf{s}(0, -e) = (0, -1, \dots, 1-e)$.

Suppose that $\mathbf{s} := \mathbf{s}(i, N)$ is a segment and let $\alpha = |\mathbf{s}| \in Q_+$. Define the corresponding *segment module* $M(\mathbf{s}) := \mathcal{O} \cdot m(\mathbf{s})$ to be the graded R_α -module which is the free $\mathcal{O}$ -module of rank one on the generator $m(\mathbf{s})$ of degree 0 with action

$$e(\mathbf{i})m(\mathbf{s}) = \delta_{\mathbf{i}, \mathbf{s}}m(\mathbf{s}), \quad \psi_r m(\mathbf{s}) = 0 \quad \text{and} \quad y_t m(\mathbf{s}) = 0$$

for all admissible $\mathbf{i}$, r and t . Equivalently, $M(\mathbf{s}) = R_\alpha / K(\mathbf{s})$, where $K(\mathbf{s})$ is the left ideal of R_α generated by the elements $e(\mathbf{i}) - \delta_{\mathbf{i}, \mathbf{s}}$, ψ_r , and y_t , for all admissible $\mathbf{i}$, r and t .

Let $\vec{s} = (\mathbf{s}(1), \dots, \mathbf{s}(n))$ be an ordered tuple of segments. Set $\alpha_r := |\mathbf{s}(r)|$, and let $\lambda_r := \text{ht}(\alpha_r)$ be the length of the segment $\mathbf{s}(r)$, for $r = 1, \dots, n$. Also set $\alpha = \alpha_1 + \dots + \alpha_n$ and $d := \text{ht}(\alpha)$. Note that $(\lambda_1, \dots, \lambda_n)$ is a composition of d . Define the *permutation module*

$$M(\vec{s}) = M(\mathbf{s}(1), \dots, \mathbf{s}(n)) := M(\mathbf{s}(1)) \circ \dots \circ M(\mathbf{s}(n)).$$

This is the graded R_α -module generated by the vector

$$m(\vec{s}) := 1 \otimes m(\mathbf{s}(1)) \otimes \dots \otimes m(\mathbf{s}(n)) \quad (3.20)$$

in $\text{Ind}_{\alpha_1, \dots, \alpha_n}^\alpha M(\mathbf{s}(1)) \boxtimes \dots \boxtimes M(\mathbf{s}(n))$. Let $\mathfrak{S}_{\vec{s}}$ be the parabolic subgroup $\mathfrak{S}_{\lambda_1} \times \dots \times \mathfrak{S}_{\lambda_n}$ of $\mathfrak{S}_d$. Define

$$\mathbf{j}(\vec{s}) := \mathbf{s}(1) \dots \mathbf{s}(n) \in I^\alpha \quad (3.21)$$

where the product on the right hand side is the concatenation. Now let $K(\vec{s})$ to be the left ideal of R_α generated by

$$\{e(\mathbf{i}) - \delta_{\mathbf{i}, \mathbf{j}(\vec{s})}, y_r, \psi_t \mid \mathbf{i} \in I^\alpha, 1 \leq r \leq d, 1 \leq t < d \text{ such that } s_t \in \mathfrak{S}_{\vec{s}}\}.$$

Then we have:

$$M(\vec{s}) \cong R_\alpha / K(\vec{s}). \quad (3.22)$$

Under this isomorphism $m(\vec{s})$ is identified with $1 + K(\vec{s})$. With the notation as above, we have as an immediate consequence of the Basis Theorem 3.18:

Theorem 3.23. *Let $\mathcal{D}_{\vec{s}}$ be the set of the shortest length left coset representatives of $\mathfrak{S}_{\vec{s}}$ in $\mathfrak{S}_d$. Then $\{\psi_w m(\vec{s}) \mid w \in \mathcal{D}_{\vec{s}}\}$ is an $\mathcal{O}$ -basis of $M(\vec{s})$. Moreover each basis element $\psi_w m(\vec{s})$ is homogeneous of degree equal to the degree of the element $\psi_w e(\mathbf{j}(\vec{s})) \in R_\alpha$, and $\psi_w m(\vec{s}) \in e(w \cdot \mathbf{j}(\vec{s}))M(\vec{s})$.*

4. BLOCK INTERTWINERS

Throughout this section we assume that $e > 0$. Recall from (2.2) that $\delta \in Q_+$ is the null-root. We fix $i \in I$ and a *composition* $\lambda = (\lambda_1, \dots, \lambda_n)$ of k . Define

$$\vec{s}(i, \lambda) := (\mathbf{s}(i, e\lambda_1), \dots, \mathbf{s}(i, e\lambda_n))$$

the tuple of segment of lengths $e\lambda_1, \dots, e\lambda_n$, all starting at i . We consider the corresponding permutation module

$$M(i, \lambda) := M(\vec{s}(i, \lambda))$$

over the algebra $R_{k\delta}$ as in section 3.6. Let

$$\mathbf{j} = (j_1, \dots, j_{ke}) := \mathbf{j}(\vec{s}(i, \lambda)) \in I^{k\delta}$$

as defined in (3.21). We have $\mathbf{j} = \mathbf{s}(i, ke)$. Finally, let the corresponding idempotent be

$$e(i, \lambda) := e(\mathbf{j}(\vec{s}(i, \lambda))) \in R_{k\delta}$$

and

$$m(i, \lambda) := m(\vec{s}(i, \lambda)) \in M(i, \lambda),$$

the generator of $M(\mathbf{s})$ as in (3.20).

4.1. The elements σ . We consider the element w_r of the symmetric group $\mathfrak{S}_{ke}$ defined as the product of transpositions

$$w_r := \prod_{a=re-e+1}^{re} (a, a+e) \quad (1 \leq r < k). \quad (4.1)$$

Informally, w_r permutes the r th “ e -block” and the $(r+1)$ st “ e -block”. If we write $w_r = w'_r s_{re}$ then $\ell(w_r) = \ell(w'_r) + 1$.

Define

$$\sigma_r := \psi_{w_r} e(i, \lambda) \in R_{k\delta} \quad (1 \leq r < k). \quad (4.2)$$

Note by Lemma 3.17 that w_r and w'_r are fully commutative elements so the elements ψ_{w_r} and $\psi_{w'_r}$ of $R_{k\delta}$ do not depend on the choice of preferred reduced decompositions for these permutations. Furthermore, $\psi_{w_r} = \psi_{w'_r} \psi_{re}$.

To prove the results in this section we will use the graphical representation of elements of $R_{k\delta}$ and $M(i, \lambda) = R_{k\delta}/K(\vec{s}(i, \lambda)) = R_{k\delta}m(i, \lambda)$ following

[16]. In fact, the diagram used to represent an element $he(i, \lambda) \in R_{k\delta}$ in [16], here will represent the element $hv \in M(i, \lambda)$, for some $v \in e(i, \lambda)M(i, \lambda)$. Of course, the element v needs to be specified before this makes sense. For example, if $v = m(i, \lambda)$, then

$$m(i, \lambda) = \begin{array}{c} j_1 \quad j_2 \quad \dots \quad j_{ke} \\ | \quad | \quad \dots \quad | \\ \vdots \quad \vdots \quad \dots \quad \vdots \end{array}, \quad \psi_r m(i, \lambda) = \begin{array}{c} j_1 \quad j_{r-1} j_r j_{r+1} \quad \dots \quad j_{ke} \\ | \quad | \quad \times \quad | \quad \dots \quad | \\ \vdots \quad \vdots \quad \times \quad \vdots \quad \dots \quad \vdots \end{array},$$

and

$$y_s m(i, \lambda) = \begin{array}{c} j_1 \quad j_{s-1} j_s \quad j_s \quad \dots \quad j_{ke} \\ | \quad | \quad \bullet \quad | \quad \dots \quad | \\ \vdots \quad \vdots \quad \bullet \quad \vdots \quad \dots \quad \vdots \end{array},$$

where $1 \leq r < d$ and $1 \leq s \leq d$. Also, setting $r' = (r-1)e$, $r'' = (r+1)e+1$, we have

$$\sigma_r m(i, \lambda) = \begin{array}{c} j_1 \quad j_{r'} \quad i \quad i+1 \quad \dots \quad i-1 \quad i \quad i+1 \quad \dots \quad i-1 \quad j_{r''} \quad j_{ke} \\ | \quad | \quad \times \quad \times \quad \times \quad \times \quad \times \quad \times \quad \times \quad | \quad | \\ \vdots \quad \vdots \quad \times \quad \times \quad \times \quad \times \quad \times \quad \times \quad \times \quad \vdots \quad \vdots \end{array} \quad (4.3)$$

We will colour the strings of the diagrams to improve readability, but these colours will have no mathematical meaning (and will not be distinguishable in black and white!).

4.2. The block permutation subspace. Consider the *block permutation subspace*

$$T(i, \lambda) := \mathcal{O}\text{-span}\{\sigma_{r_1} \sigma_{r_2} \dots \sigma_{r_a} m(i, \lambda) \mid 1 \leq r_1, \dots, r_a < k\}.$$

It is not hard to see using Theorem 3.23 that

$$T(i, \lambda) = e(i, \lambda)M(i, \lambda).$$

It is easy to see that $\deg(\sigma_r e(i, \lambda)) = 0$. Therefore,

$$T(i, \lambda) \subseteq M(i, \lambda)_0,$$

the degree zero component of $M(i, \lambda)$.

Lemma 4.4. *Suppose that $1 \leq s, t \leq ke$ with $t \not\equiv 0 \pmod{e}$. Then the elements y_s and ψ_t act as zero on $T(i, \lambda)$.*

Proof. Let $v \in T(i, \lambda)$. We have

$$y_s v \in e(i, \lambda)M(i, \lambda) = T(i, \lambda) \subseteq M(i, \lambda)_0.$$

On the other hand, $\deg(y_s v) = \deg(y_s) + \deg(v) = 2$. Hence $y_s v = 0$. Moreover, $\psi_t v \in e(s_t \cdot \mathbf{j})M(i, \lambda) = 0$, the last equality holding by Theorem 3.23. $\square$

4.3. Quadratic relation. We want to study relations satisfied by the elements σ_r acting on $T(i, \lambda)$. Our main goal is to show that the elements $\tau_r := \sigma_r + 1$ satisfy the Coxeter relations on $T(i, \lambda)$. We begin with the quadratic relations.

Lemma 4.5. *Suppose that $1 \leq r < k$ and $v \in T(i, \lambda)$. Then $\psi_{re}\sigma_r v = -2\psi_{re}v$. Equivalently, in terms of diagrams we have*

where $r' = (r-1)e$, $r'' = (r+1)e+1$ and $\mathbf{i} = \mathbf{s}(i, ke)$.


Proof. For typographical convenience, we only consider the case where $r = 1$ and $i = 0$. We first treat the case $e = 2$ which is exceptional because in this case the quiver Γ is not simply laced. Using the relation (3.9), and then (3.7) and (3.8), we have:

as required.


Now suppose that $e > 2$. To start, using (3.9) we see that $\psi_e \sigma_1 v$ equals

Let D_1 be the first diagram and let D_2 be the second diagram. To complete the proof, we show that $D_1 = \psi_e v$ and $D_2 = -\psi_e v$. In fact, the two equalities are proved similarly, so we give details only for the first one. Using (3.7), we see that


The second summand is zero as $y_1 v = 0$ by Lemma 4.4. Applying the braid relations (3.10) to the first summand, we get that D_1 equals




Using the braid relations to pull the second 0-string through shows that the first summand equals




showing that this element is zero since $\psi_{e+1} v = 0$, by Lemma 4.4. Applying the braid relations to the second summand, we get




As before, using the braid relations to pull the second 1-string to the top of the first diagram shows that the first summand is zero. So, by (3.9),



The argument so far has straightened the first three strings in the diagram. Continuing in this way straightens the first $e - 1$ strings so that



Now applying the braid relation for the last time shows that D_1 equals



as required. □

Recall from section 4.1 that $\psi_{w_r} = \psi_{w'_r} \psi_{re}$.

Corollary 4.6. *Suppose that $1 \leq r < k$ and $v \in T(i, \lambda)$. Then $\sigma_r^2 v = -2\sigma_r v$.*

Proof. Using Lemma 4.5, we get

$$\sigma_r^2 v = \psi_{w'_r} \psi_{re} \sigma_r v = -2\psi_{w'_r} \psi_{re} v = -2\psi_{w_r} v = -2\sigma_r v,$$

as desired. $\square$

4.4. Braid relations. This section is dedicated to the proof of the following

Theorem 4.7. *Suppose that $1 \leq r < k$ and $v \in T(i, \lambda)$. Then*

$$(\sigma_r \sigma_{r+1} \sigma_r - \sigma_{r+1} \sigma_r \sigma_{r+1} - \sigma_r + \sigma_{r+1})v = 0.$$

In the proof, for typographical reasons, we assume that $i = 0$ and $k = 3$ (this corresponds to ignoring vertical strings to the left and to the right of the relation we are interested in). As in the Lemma 4.5, all diagrams represent elements of $T(i, \lambda)$ obtained by applying the corresponding elements of $R_{k\delta}$ to a given $v \in T(i, \lambda)$.

First, we need three technical lemmas.

Lemma 4.8. *Suppose that $e > 2$, $k = 3$, $i = 0$, and $v \in T(i, \lambda)$. Then:*

$$\sigma_1 v = \text{[Diagram 1]} ,$$

and

$$\sigma_2 v = \text{[Diagram 2]} .$$

Proof. We prove only the first identity for σ_1 as the proof of the second one is almost identical. Let D_1 be the first diagram on the right hand side of the first equality. Using more strings for clarity of exposition,

$$D_1 = \text{[Diagram 3]} .$$

Pulling the rightmost 0-string past the ${}^1 \times {}^1$ -crossing immediately to its right gives zero because $\psi_{2e+1} v = 0$ by Lemma 4.4. Here, and in similar situations

below, we will omit such terms which arise when applying the braid relations (3.10). This observation shows that

$$D_1 = \begin{array}{c} \begin{array}{cccccccccccccccccccccccccccccccccccccccc} 0 & 1 & 2 & & -3 & -2 & -1 & 0 & 1 & 2 & & -3 & -2 & -1 & 0 & 1 & 2 & & -3 & -2 & -1 & 0 & 1 & 2 & & -3 & -2 & -1 \end{array} \\ \text{[Braid diagram with blue and red strings]} \end{array} \\ = \begin{array}{c} \begin{array}{cccccccccccccccccccccccccccccccccccccccc} 0 & 1 & 2 & & -3 & -2 & -1 & 0 & 1 & 2 & & -3 & -2 & -1 & 0 & 1 & 2 & & -3 & -2 & -1 & 0 & 1 & 2 & & -3 & -2 & -1 \end{array} \\ \text{[Braid diagram with blue and red strings]} \end{array}$$

where for the second equality we pulled the rightmost 1-string past the ${}^2\bowtie^2$ -crossing. Continuing in this way and pulling the right most $(i-1)$ -string past its neighbouring ${}^i\bowtie^i$ -crossing, for $3 \leq i < e-1$, shows that

$$D_1 = \begin{array}{c} \begin{array}{cccccccccccccccccccccccccccccccccccccccc} 0 & 1 & 2 & & -3 & -2 & -1 & 0 & 1 & 2 & & -3 & -2 & -1 & 0 & & & & & -5 & -4 & -3 & -2 & -1 \end{array} \\ \text{[Braid diagram with blue strings]} \end{array} .$$

Another application of the braid relation (3.10) yields

$$D_1 = \begin{array}{c} \begin{array}{cccccccccccccccccccccccccccccccccccccccc} 0 & 1 & 2 & & -3 & -2 & -1 & 0 & 1 & 2 & & -3 & -2 & -1 & 0 & & & & & -5 & -4 & -3 & -2 & -1 \end{array} \\ \text{[Braid diagram with blue and red strings]} \end{array} .$$

Applying (3.6) we can straighten the rightmost $-3, -4, \dots, -1$ strings completely and then pull the next $e+1$ strings to the right to give

$$D_1 = \begin{array}{c} \begin{array}{cccccccccccccccccccccccccccccccccccccccc} 0 & 1 & 2 & & -3 & -2 & -1 & 0 & 1 & 2 & 3 & & -2 & -1 & 0 & 1 & 2 & & -3 & -2 & -1 \end{array} \\ \text{[Braid diagram with blue strings]} \end{array} .$$

Now applying the quadratic relation (3.9) to the rightmost pair of $(-1, 0)$ -strings, using Lemma 4.4, and then applying the relation (3.8), gives

$$D_1 = \text{Diagram 1} = \text{Diagram 2}$$

Repeating the same argument another $e - 2$ times shows that

$$D_1 = \text{Diagram 3}$$

A final application of (3.9) and (3.8) now shows that $D_1 = \sigma_1 v$ completing the proof. $\square$

Lemma 4.9. Suppose that $k = 3$, $i = 0$, $v \in T(i, \lambda)$ and let

$$E_1 = \text{Diagram 4}, \quad E'_1 = \text{Diagram 5},$$

$$E_2 = \text{Diagram 6}, \quad E'_2 = \text{Diagram 7}$$

Then $E_1 = \sigma_1 v + E'_1$ and $E_2 = \sigma_2 v + E'_2$.

Proof. Both identities are proved similarly, so we consider only the first one. First consider the exceptional case $e = 2$. Then we have to show that

$$\text{Diagram 8} = \sigma_1 v + \text{Diagram 9} \quad (4.10)$$

Applying the braid relation (3.10), (the first line of) the quadratic relation (3.9) and Lemma 4.4, shows that

$$\text{Diagram 1} = \text{Diagram 2} + \text{Diagram 3}.$$

Applying (3.10) twice to the first summand and (3.8) to the second summand gives

$$\text{Diagram 1} = \text{Diagram 2} + \text{Diagram 3}.$$

The relations (3.10), (3.8) and Lemma 4.4 show that the first summand above equals the second summand on the right hand side of (4.10) and the second summand above equals $\sigma_1 v$.

Now consider the case when $e > 2$. By (3.10), E_1 is equal to

$$\text{Diagram 1} + \text{Diagram 2}.$$

By Lemma 4.8 the second summand is equal to $\sigma_1 v$. Using the braid relations again, the first summand is equal to

$$\text{Diagram 1} = E'_1 + \text{Diagram 2}.$$

Using the braid relations to pull the rightmost -2 -string in the second summand above to the right and observing that the error term of the braid relation equals zero by (3.9), shows that the second summand equals

$$D = \text{Diagram 1} = 0,$$

where the last equality follows because $\psi_{3e-1} v = 0$ in view of Lemma 4.4. Therefore, $E_1 = \sigma_1 v + E'_1$ as claimed. $\square$

Lemma 4.11. *Suppose that $i \neq 0, -1$ and $v \in T(i, \lambda)$. Then*

$$\text{Diagram 1} = \text{Diagram 2}.$$

Proof. Let D be the left hand diagram. Then, using the braid relations,

$$D = \begin{array}{c} \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} \\ + \\ \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array} \end{array} .$$

The diagrams consist of blue and red strings crossing in a complex braid-like pattern. The top and bottom rows of each diagram are labeled with sequences of numbers: $0, i-1, i, i+1, -1, 0, i-1, i, i+1, -1, 0, i-1, i, i+1, -1$.

Let the first summand of D be D_1 and the second one be D_2 . Then by the braid relations, we have

$$D_1 = \begin{array}{c} \begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \end{array} \\ + \\ \begin{array}{c} \text{Diagram 7} \\ \text{Diagram 8} \end{array} \end{array} .$$


The diagrams are similar to the previous ones, with blue and red strings and the same top/bottom labels.

The first summand is zero because we can use (3.10) to pull the rightmost $(i-1)$ -string to the top of the diagram and then use the fact that $\psi_{2e+i}v = 0$ by Lemma 4.4. The second summand is zero by because $\textcolor{blue}{\bowtie}^i = 0$ by (3.9). Hence, $D_1 = 0$. Now consider D_2 . Using the braid relations to pull the middle i -string in D_2 to the right, D_2 is equal to the diagram on the right hand side of the formula in the statement of the lemma plus the following error term


$$\begin{array}{c} \text{Diagram 9} \\ \text{Diagram 10} \end{array} ,$$

The diagrams show the error term, with blue and red strings and the same top/bottom labels.

which using the (error term free) braid relations, equals




By the braid relations again, this equals



The first summand is zero since $\psi_i v = 0$ by Lemma 4.4. The second term is zero because of the quadratic relation $\textcolor{blue}{\bowtie}^i = 0$. The proof of the lemma is complete. $\square$

We can now prove Theorem 4.7.

Proof of Theorem 4.7. Writing $\sigma_1 \sigma_2 \sigma_1 v$ in terms of diagrams and using Lemma 4.9 we have



Hence, applying Lemma 4.11 ($e - 1$) times,

$$\begin{aligned}
 \sigma_1 \sigma_2 \sigma_1 v &= \sigma_1 v + \\
 &\quad \text{[Diagram: A sequence of blue lines representing a permutation. The top row of labels is 0 1 -2 -1 0 1 -2 -1 0 1 -2 -1. The lines cross in a pattern that corresponds to the permutation $\sigma_1 \sigma_2 \sigma_1$.]} \\
 &= \cdots = \sigma_1 v + \\
 &\quad \text{[Diagram: A sequence of blue lines representing a permutation. The top row of labels is 0 1 -2 -1 1 0 -2 -1 1 0 -2 -1. The lines cross in a pattern that corresponds to the permutation $\sigma_2 \sigma_1 \sigma_2$ applied to the previous diagram.]} \\
 &= \sigma_1 v - \sigma_2 v + \\
 &\quad \text{[Diagram: A sequence of blue lines representing a permutation. The top row of labels is 0 1 -2 -1 0 1 -2 -1 0 1 -2 -1. The lines cross in a pattern that corresponds to the permutation σ_2 applied to the previous diagram.]} ,
 \end{aligned}$$

where the last equality follows using the identity $E_2 = \sigma_2 v + E'_2$ from Lemma 4.9. The diagram in the last equation is equal to $\sigma_2 \sigma_1 \sigma_2 v$, so this completes the proof of Theorem 4.7. $\square$

4.5. The elements τ . Let $1 \leq r < k$. Recall from the beginning of the section that $\sigma_r = \psi_{w_r} e(i, \lambda) \in R_{k\delta}$. Define

$$\tau_r = (\sigma_r + 1)e(i, \lambda) = \sigma_r + e(i, \lambda) = (\psi_{w_r} + 1)e(i, \lambda).$$

Quite remarkably, as we now show, the elements $\tau_1, \dots, \tau_{k-1}$ satisfy the usual Coxeter relations for the symmetric group $\mathfrak{S}_k$ when they act on the block permutation space $T(i, \lambda)$. Let $\mathfrak{S}_\lambda = \mathfrak{S}_{\lambda_1} \times \dots \times \mathfrak{S}_{\lambda_n}$ be the parabolic subgroup of $\mathfrak{S}_k$ indexed by λ , $\mathcal{O}_\lambda$ the trivial representation of $\mathfrak{S}_\lambda$, and $\mathcal{D}_\lambda$ the set of the minimal length left coset representatives of $\mathfrak{S}_\lambda$ in $\mathfrak{S}_k$.

Theorem 4.12. *Suppose that $1 \leq r, s < k$ and $v \in T(i, \lambda)$. Then*

- (i) $\tau_r^2 v = v$.
- (ii) If $|r - s| > 1$ then $\tau_r \tau_s v = \tau_s \tau_r v$.
- (iii) If $r < k - 1$ then $\tau_r \tau_{r+1} \tau_r v = \tau_{r+1} \tau_r \tau_{r+1} v$.

Consequently, $\mathfrak{S}_k$ acts on $T(i, \lambda)$, and the elements $\tau_u m(i, \lambda)$ for $u \in \mathfrak{S}_k$ are well-defined. Finally, $T(i, \lambda) \cong \text{ind}_{\mathcal{O}_{\mathfrak{S}_\lambda}}^{\mathcal{O}_{\mathfrak{S}_k}} \mathcal{O}_\lambda$ as $\mathcal{O}_{\mathfrak{S}_k}$ -modules, and $T(i, \lambda)$ has $\mathcal{O}$ -basis $\{\tau_u m(i, \lambda) \mid u \in \mathcal{D}_\lambda\}$.

Proof. Part (i) comes from Corollary 4.6. Part (ii) follows directly from the definition of σ_r . For part (iii), by definition

$$\begin{aligned}
 \tau_r \tau_{r+1} \tau_r v &= (\sigma_r + 1)(\sigma_{r+1} + 1)(\sigma_r + 1)v \\
 &= (\sigma_r \sigma_{r+1} \sigma_r + \sigma_r \sigma_{r+1} + \sigma_{r+1} \sigma_r + \sigma_r^2 + 2\sigma_r + \sigma_{r+1} + 1)v.
 \end{aligned}$$

Theorem 4.7 and Corollary 4.6 now imply (iii). So we obtain the action of the symmetric group $\mathfrak{S}_k$ on $T(i, \lambda)$ with Coxeter generators s_r acting as τ_r for all $r = 1, \dots, k - 1$.

For the final statement of the proposition, consider the parabolic subgroup $\mathfrak{S}_\lambda \leq \mathfrak{S}_k$ generated by

$$\{s_r \mid r \neq \lambda_1 + \cdots + \lambda_a \text{ for all } a = 1, \dots, n\}.$$

Note from the definition of $T(i, \lambda)$ that $T(i, \lambda)$ is an $\mathcal{O}$ -span of all elements of the form $\tau_{r_1} \cdots \tau_{r_a} m(i, \lambda)$. Moreover, $\mathcal{O} \cdot m(i, \lambda) \cong \mathcal{O}_\lambda$ is the trivial module of $\mathfrak{S}_\lambda$ because if $s_r \in \mathfrak{S}_\lambda$ then $\tau_r m(i, \lambda) = m(i, \lambda)$ since $\sigma_r m(i, \lambda) = 0$ by (3.20). So we have a surjective homomorphism from $\text{ind}_{\mathcal{O}_{\mathfrak{S}_\lambda}}^{\mathcal{O}_{\mathfrak{S}_k}} \mathcal{O}_\lambda$ onto $T(i, \lambda)$, which sends the natural cyclic generator of $\text{ind}_{\mathcal{O}_{\mathfrak{S}_\lambda}}^{\mathcal{O}_{\mathfrak{S}_k}} \mathcal{O}_\lambda$ onto $m(i, \lambda)$. The injectivity of this map follows from Theorem 3.23, which describes an $\mathcal{O}$ -basis for $M(i, \lambda)$, together with the observation that the transition matrix for the change of basis from the products of the τ_r to the corresponding products of the σ_r is unitriangular. $\square$


5. HOMOGENEOUS GARNIR RELATIONS

In this section we define universal graded (row) Specht modules S^μ for R_α by generators and relations, see Definition 5.9. This definition will be justified in Theorem 6.23 when we show that these universal graded Specht modules are isomorphic to the usual graded Specht modules from [6, 11].

5.1. Row Garnir tableaux. The definitions here differ slightly from those given in [6] but match those in [21]. Let $A = (a, b, m)$ be a node of $\mu \in \mathcal{P}^\kappa$. Then A is a (row) *Garnir node* if $(a+1, b, m)$ is also a node of μ . The (row) *A-Garnir belt* $\mathbf{B}^A$ is the set of nodes

$$\mathbf{B}^A = \{ (a, c, m) \in \mu \mid b \leq c \leq \mu_a^{(m)} \} \cup \{ (a+1, c, m) \in \mu \mid 1 \leq c \leq b \}.$$

For example, if $A = (2, 3, 2)$ then the A -Garnir belt $\mathbf{B}^A$ for $\mu = ((1), (7, 7, 4, 1))$ is highlighted below:



The (row) *A-Garnir tableau* is the μ -tableaux $\mathbf{G}^A$ defined as follows. Let $u = \mathbf{T}^\mu(a, b, m)$ and $v = \mathbf{T}^\mu(a+1, b, m)$. Now insert the numbers $u, u+1, \dots, v$ into the nodes of the Garnir belt going from left bottom to top right, and the other numbers into the same positions as in $\mathbf{T}^\mu$. Continuing the previous example, $u = 11, v = 18$, and $\mathbf{T}^\mu$ and the $(2, 3, 2)$ -Garnir tableau are:

$$\begin{array}{c} \boxed{1} \\ \mathbf{T}^\mu = \begin{array}{|c|c|c|c|c|c|c|c|} \hline 2 & 3 & 4 & 5 & 6 & 7 & 8 & \\ \hline 9 & 10 & 11 & 12 & 13 & 14 & 15 & \\ \hline 16 & 17 & 18 & 19 & & & & \\ \hline 20 & & & & & & & \\ \hline \end{array}, \quad \mathbf{G}^A = \begin{array}{|c|c|c|c|c|c|c|c|} \hline 2 & 3 & 4 & 5 & 6 & 7 & 8 & \\ \hline 9 & 10 & 14 & 15 & 16 & 17 & 18 & \\ \hline 11 & 12 & 13 & 19 & & & & \\ \hline 20 & & & & & & & \\ \hline \end{array} \end{array}$$

Lemma 5.1. *Suppose that $\mu \in \mathcal{P}^\kappa$, A is a Garnir node of μ , and $\mathbf{S} \in \text{St}(\mu)$. If $\mathbf{S} \triangleright \mathbf{G}^A$ then $\mathbf{S}$ agrees with $\mathbf{T}^\mu$ outside the A -Garnir belt.*

Proof. This follows from [6, Lemma 3.9] and Lemma 2.19. $\square$

The importance of the Garnir tableaux comes from the following:

Lemma 5.2. *Suppose that $\mu \in \mathcal{P}_d^\kappa$ and that $\mathbf{T}$ is a row-strict μ -tableau which is not standard. Then there exists a Garnir tableau $\mathbf{G} = \mathbf{G}^A$, for $A \in \mu$ a row Garnir node, and $w \in S_d$ such that $\mathbf{T} = w\mathbf{G}$ and $\ell(w^{\mathbf{T}}) = \ell(w^{\mathbf{G}}) + \ell(w)$.*

Proof. If $l = 1$ this is [21, Lemma 3.14], and the general case follows easily from the case $l = 1$. $\square$

5.2. Bricks. Fix $\mu \in \mathcal{P}_d^\kappa$ and a Garnir node $A = (a, b, m) \in \mu$. A (row A -)brick is a set of e successive nodes in the same row

$$\{(c, d, m), (c, d + 1, m), \dots, (c, d + e - 1, m)\} \subseteq \mathbf{B}^A$$


such that $\text{res}(c, d, m) = \text{res } A$. Note that $\mathbf{B}^A$ is a disjoint union of the bricks that it contains together with less than e nodes at the end of row a which are not contained in a brick and less than e nodes at the beginning of row $a + 1$ which are not contained in a brick.

Let $k = k^A$ be the number of bricks in $\mathbf{B}^A$. We label the bricks

$$B_1^A, B_2^A, \dots, B_k^A$$

going from left to right along row $a + 1$ and then from left to right along row a of $\mathbf{G}^A$ as in the example above. Of course, it might happen that $\mathbf{B}^A$ does not contain any bricks (this is always true if $e = 0$), in which case $k = 0$.

For example, the following diagram shows the bricks in the $(2, 3, 2)$ -Garnir belt of $\mu = ((1), (7, 7, 4, 1))$ when $e = 2$:



Note that $k = 3$, there are two bricks B_2^A, B_3^A in row 2 and one brick B_1^A in row 3 of the second component. Further, $(3, 1, 2)$ and $(2, 7, 2)$ are the only nodes in the $(2, 3, 2)$ -Garnir belt of $\mathbf{G}^A$ which are not contained in a brick.

Assume now that $k > 0$ and let $n = n^A$ be the smallest entry in $\mathbf{G}^A$ which is contained in a brick in $\mathbf{B}^A$. In the example above, $n = 12$. Extending (4.1), define

$$w_r^A = \prod_{a=n+re-e}^{n+re-1} (a, a+e) \in \mathfrak{S}_d \quad (1 \leq r < k). \quad (5.3)$$



Informally, w_r^A swaps the bricks B_r^A and B_{r+1}^A . The elements $w_1^A, w_2^A, \dots, w_{k-1}^A$ are the Coxeter generators of the symmetric group

$$\mathfrak{S}^A := \langle w_1^A, w_2^A, \dots, w_{k-1}^A \rangle \cong \mathfrak{S}_k.$$


We call $\mathfrak{S}^A$ the (row) *brick permutation group*. By convention, $\mathfrak{S}^A$ is the trivial group if $k = 0$.

Define $f = f^A$ to be the number of A -bricks in row a of the Garnir belt $\mathbf{B}^A$. Finally, let $\mathscr{D}^A$ be the set of minimal length left coset representations of $\mathfrak{S}_f \times \mathfrak{S}_{k-f}$ in $\mathfrak{S}^A \cong \mathfrak{S}_k$. Note that by definition $\mathfrak{S}^A$ is a subgroup of $\mathfrak{S}_d$, so $\mathscr{D}^A$ is a subset of $\mathfrak{S}_d$ and, in particular, its elements act on μ -tableaux. Note that

Continuing the example above, T^A is the tableau



and



The circles in these diagrams correspond to the bricks in the Garnir belt.

The degree statement in the following lemma is what will guarantee the homogeneity of our Garnir relations. This result is implicit in the proof of [6, Proposition 3.16].

Lemma 5.5. *Suppose that $\mu \in \mathcal{P}_d^\kappa$ and $A \in \mu$ is a Garnir node. Then*

$$\text{Gar}^A \setminus \{\mathbf{G}^A\} = \{\mathbf{T} \in \text{St}(\mu) \mid \mathbf{T} \supseteq \mathbf{G}^A \text{ and } \mathbf{i}(\mathbf{T}) = \mathbf{i}^A\}.$$

Moreover, $\deg(\mathbf{T}) = \deg(\mathbf{G}^A)$ for all $\mathbf{T} \in \text{Gar}^A$.

5.3. The row permutation modules M^μ . Let $\mu \in \mathcal{P}_\alpha^\kappa$ be a multipartition with (non-empty) rows $R_1, \dots, R_g$ counted from top to bottom. If a row R_a has length N and the leftmost node of R_a has residue i we associate the segment $\mathbf{r}(a) := s(i, N)$ to R_a . Let $\vec{\mathbf{r}} = (\mathbf{r}(1), \dots, \mathbf{r}(g))$, and, recalling the definitions from section 3.6, put

$$M^\mu = M^\mu(\mathcal{O}) := M(\vec{\mathbf{r}}) \langle \deg \mathbf{T}^\mu \rangle.$$

Note the degree shift by $\deg \mathbf{T}^\mu$, the significance of which is explained by Theorem 6.23 below. The module M^μ is generated by the vector $m^\mu := m(\vec{\mathbf{r}})$ of degree $\deg \mathbf{T}^\mu$. Recalling (3.16), for any μ -tableau $\mathbf{T}$ we define

$$m^\mathbf{T} := \psi^\mathbf{T} m^\mu.$$

The following is a special case of Theorem 3.23:

Theorem 5.6. *Suppose that $\alpha \in Q_+$ and $\mu \in \mathcal{P}_\alpha^\kappa$. Then*

$$\{m^\mathbf{T} \mid \mathbf{T} \text{ is a row-strict } \mu\text{-tableau}\}$$

is an $\mathcal{O}$ -basis of M^μ .

5.4. Universal row Specht modules S^μ . Fix a Garnir node $A \in \mu$, and let $\mathfrak{S}^A$ be the corresponding block permutation group with generators $w_1^A, \dots, w_{k-1}^A$ as defined in section 5.2. Using the notation of (5.3), we define

$$\sigma_r^A := \psi_{w_r^A} e(\mathbf{i}^A) \quad \text{and} \quad \tau_r^A := (\sigma_r^A + 1) e(\mathbf{i}^A), \quad (5.7)$$

cf. section 4.5. Any element $u \in \mathfrak{S}^A$ can be written as a reduced product $u = w_{r_1}^A \dots w_{r_a}^A$ of simple generators $w_1^A, \dots, w_{k-1}^A$ of $\mathfrak{S}^A$. In general, the elements τ_r^A do not have to satisfy Coxeter relations. However, if u is fully commutative then the element

$$\tau_u^A := \tau_{r_1}^A \dots \tau_{r_a}^A$$

is well-defined, since τ_r^A and τ_s^A commute for $|r - s| > 1$. In particular, we have well-defined elements

$$\{\tau_u^A \mid u \in \mathcal{D}^A\}.$$

(As operators on the brick permutation space $T^{\mu, A}$, defined below, the elements τ_r^A do satisfy Coxeter relations, see Theorem 5.11(ii).)

Recall from (5.4) that Gar^A is the set of row-strict tableaux obtained from the tableau $\mathbf{T}^A$ by acting with the elements of $\mathcal{D}^A$. Note that for any $\mathbf{S} \in \text{Gar}^A$, we can write $w^{\mathbf{S}} = u^{\mathbf{S}} w^{\mathbf{T}^A}$ so that $\ell(w^{\mathbf{S}}) = \ell(u^{\mathbf{S}}) + \ell(w^{\mathbf{T}^A})$ and $u^{\mathbf{S}} \in \mathcal{D}^A$. Moreover, in view of Lemma 3.17, all elements $w^{\mathbf{S}}, u^{\mathbf{S}}$, and $w^{\mathbf{T}^A}$ are fully commutative so the elements $\psi_{u^{\mathbf{S}}}$, $\psi^{\mathbf{T}^A}$ and $\psi^{\mathbf{S}} = \psi_{u^{\mathbf{S}}} \psi^{\mathbf{T}^A}$ are all independent of the choice of preferred reduced decomposition. Set

$$m^A := m^{\mathbf{T}^A} = \psi^{\mathbf{T}^A} m^{\mu} \in M^{\mu}.$$

Definition 5.8. Suppose that $\mu \in \mathcal{P}_{\alpha}^{\kappa}$ and $A \in \mu$ is a Garnir node. The (row) Garnir element is

$$g^A := \sum_{u \in \mathcal{D}^A} \tau_u^A \psi^{\mathbf{T}^A} \in R_{\alpha}.$$

In the module M^{μ} we have

$$g^A m^{\mu} = \sum_{u \in \mathcal{D}^A} \tau_u^A m^A.$$

By Lemma 5.5 all of the summands on the right hand side have the same degree. Finally, if $\mathcal{D}^A = \{1\}$, we have $\mathbf{G}^A = \mathbf{T}^A$ and $g^A = \psi^{\mathbf{G}^A}$.

Definition 5.9. Let $\alpha \in Q_+$, $d = \text{ht}(\alpha)$, and $\mu \in \mathcal{P}_{\alpha}^{\kappa}$. Define the *universal graded (row) Specht module* $S^{\mu} = S^{\mu}(\mathcal{O})$ to be the graded R_{α} -module generated by the vector z^{μ} of degree $\deg(\mathbf{T}^{\mu})$ subject only to the following relations:

- (i) $e(\mathbf{j})z^{\mu} = \delta_{\mathbf{j}, \mathbf{i}^{\mu}} z^{\mu}$ for all $\mathbf{j} \in I^{\alpha}$;
- (ii) $y_r z^{\mu} = 0$ for all $r = 1, \dots, d$;
- (iii) $\psi_r z^{\mu} = 0$ for all $r = 1, \dots, d-1$ such that $r \rightarrow_{\mathbf{T}^{\mu}} r+1$;
- (iv) (*homogeneous Garnir relations*) $g^A z^{\mu} = 0$, for all (row) Garnir nodes A in μ .

In other words, $S^{\mu} = (R_{\alpha}/J_{\alpha}^{\mu})\langle \deg(\mathbf{T}^{\mu}) \rangle$, where J_{α}^{μ} is the (homogeneous) left ideal of R_{α} generated by the elements

- (i) $e(\mathbf{j}) - \delta_{\mathbf{j}, \mathbf{i}^{\mu}}$ for all $\mathbf{j} \in I^{\alpha}$;
- (ii) y_r for all $r = 1, \dots, d$;
- (iii) ψ_r for all $r = 1, \dots, d-1$ such that $r \rightarrow_{\mathbf{T}^{\mu}} r+1$;
- (iv) g^A for all Garnir nodes $A \in \mu$.

In view of (3.22), the elements (i)-(iii) generate a left ideal K^{μ} such that $R_{\alpha}/K^{\mu} \cong M^{\mu}$. So we have a natural surjection $M^{\mu} \twoheadrightarrow S^{\mu}$, with the kernel J^{μ} of this surjection generated by the Garnir relations $g^A m^{\mu} = 0$. This surjection maps m^{μ} to z^{μ} and $J^{\mu} = J_{\alpha}^{\mu} m^{\mu}$.

Remark 5.10. Our homogeneous Garnir relations are simpler than the ones defined by Young and Garnir in that they have fewer summands. For example, if $\mathbf{G}^A$ is the only tableau in Gar^A , then the Garnir relation is simply saying that $\psi^{\mathbf{G}^A} z^{\mu} = 0$. Note that we always have $\text{Gar}^A = \{\mathbf{G}^A\}$ when $e = 0$ or $e > d$.

Our main goal is to obtain a basis for the universal Specht modules and to relate them to the usual Specht modules for cyclotomic Hecke algebras.

5.5. Row brick permutation space $T^{\mu,A}$. Continuing with the notation of the previous subsection, define the (row) brick permutation space $T^{\mu,A} \subseteq M^\mu$ to be the $\mathcal{O}$ -span of all elements of the form $\sigma_{r_1}^A \dots \sigma_{r_a}^A m^A$, cf. section 4.2.

Theorem 5.11. *Suppose that $\alpha \in Q_+$, $\mu \in \mathcal{P}_\alpha^\kappa$, $A \in \mu$ is a Garnir node and let $k = k^A$ and $f = f^A$. Then:*

- (i) $T^{\mu,A}$ is the $\mathcal{O}$ -span of all elements of the form $\tau_{r_1}^A \dots \tau_{r_a}^A m^A$. In particular, the elements $\tau_1^A, \dots, \tau_{k-1}^A$ act on $T^{\mu,A}$.
- (ii) As $\mathcal{O}$ -linear operators on $T^{\mu,A}$, the elements $\tau_1^A, \dots, \tau_{k-1}^A$ satisfy the Coxeter relations for the symmetric group $\mathfrak{S}_k$. Thus, we can consider $T^{\mu,A}$ as an $\mathcal{O}\mathfrak{S}_k$ -module.
- (iii) Let $\mathcal{O}_{(f,k-f)}$ be the trivial $\mathcal{O}\mathfrak{S}_{f,k-f}$ -module. There is an isomorphism of $\mathcal{O}\mathfrak{S}_k$ -modules

$$T^{\mu,A} \cong \text{ind}_{\mathcal{O}\mathfrak{S}_{(f,k-f)}}^{\mathcal{O}\mathfrak{S}_k} \mathcal{O}_{(f,k-f)},$$

under which $m^A \in T^{\mu,A}$ corresponds to the natural cyclic generator of the induced module on the right hand side.

- (iv) $\{\tau_u^A m^A \mid u \in \mathcal{D}^A\}$ is an $\mathcal{O}$ -basis of $T^{\mu,A}$.

Proof. Let $i = \text{res } A$, and let $n = n^A$ be, as before, the smallest entry in $\mathbf{G}^A$ which is contained in a brick in $\mathbf{B}^A$. Set $\mathbf{i} = \mathbf{i}^A := (i_1, \dots, i_d)$. For any $\mathbf{j} = (j_1, \dots, j_{ke}) \in I^{k\delta}$ define the tuple $\mathbf{j}^A = (j_1, \dots, j_d)$ where $j_t = i_t$ for $t < n$ and $t \geq n + ek$, and $j_{n+s} = j_{s+1}$ for all $s = 0, 1, \dots, ke - 1$. There is a (non-unital) embedding of algebras $\iota^A : R_{k\delta} \hookrightarrow R_\alpha$ such that

$$\iota^A : \psi_s \mapsto \psi_{s+n-1}, y_t \mapsto y_{t+n-1}, e(\mathbf{j}) \mapsto e(\mathbf{j}^A)$$

for all admissible s, t and $\mathbf{j}$. From now on we are going to suppress the notation ι^A and simply identify $R_{k\delta}$ with the subalgebra $\iota^A(R_{k\delta})$ inside R_α .

Consider the $R_{k\delta}$ -module $R_{k\delta} \cdot m^A$ generated by the vector $m^A \in M^\mu$. We claim that this module is isomorphic to the permutation module $M(i, (f, k-f))$ defined in section 4. Indeed, it is easy to check that $e(\mathbf{j})m^A = \delta_{\mathbf{j}, \mathbf{s}(i, ke)} m^A$, $y_t m^A = 0$, and $\psi_s m^A = 0$ unless $s = ef$. This shows that there is an $R_{k\delta}$ -homomorphism from $M(i, (f, k-f))$ onto $R_{k\delta} \cdot m^A$ which maps $m(\vec{\mathbf{s}}(i, (f, k-f)))$ to m^A . An application of Theorem 3.23 now implies that this homomorphism is an isomorphism. Hence, the result follows from Theorem 4.12. $\square$

Corollary 5.12. *Suppose that $\mu \in \mathcal{P}_\alpha^\kappa$, $A \in \mu$ is a Garnir node of μ , and $\mathbf{S} = u\mathbf{T}^A \in \text{Gar}^A$ for some $u \in \mathcal{D}^A$, cf. (5.4). Then*

$$m^{\mathbf{S}} = \tau_u^A m^A + \sum_{w \in \mathcal{D}^A, w \triangleleft u} c_w \tau_w^A m^A$$

for some $c_w \in \mathcal{O}$. In particular, $\{m^{\mathbf{T}} \mid \mathbf{T} \in \text{Gar}^A\}$ is an $\mathcal{O}$ -basis of $T^{\mu,A}$.

Proof. Let $u = w_{r_1}^A \dots w_{r_a}^A$ be a reduced decomposition in $\mathfrak{S}^A$. Then, using (5.7),

$$\begin{aligned} m^{\mathbf{S}} &= \psi^{\mathbf{S}} m^\mu = \psi_u \psi^{\mathbf{T}^A} m^\mu = \psi_u m^A = \sigma_{r_1}^A \dots \sigma_{r_a}^A m^A \\ &= (\tau_{r_1}^A - 1) \dots (\tau_{r_a}^A - 1) m^A, \end{aligned}$$

which implies the result in view of Theorem 5.11. $\square$

5.6. A spanning set for the universal row Specht module. Let $\alpha \in Q_+$ with $\text{ht}(\alpha) = d$, and $\mu \in \mathcal{P}_\alpha^\kappa$. Recall that $S^\mu \cong M^\mu/J^\mu$ and $z^\mu = m^\mu + J^\mu$. Also set

$$z^A := m^A + J^\mu \in S^\mu$$

for any Garnir node $A \in \mu$.

Recall from (3.16) that for each μ -tableau T we have defined the element $\psi^T \in R_\alpha$, which depends on a fixed choice of reduced decomposition of $w^T \in \mathfrak{S}_d$. Hence, we can associate to T the homogeneous element

$$v^T := \psi^T z^\mu = m^T + J^\mu \in S^\mu.$$

Lemma 5.13. *Suppose that $\mu \in \mathcal{P}_\alpha^\kappa$ and A be a Garnir node of μ . Then*

$$v^{G^A} = \sum_{T \in \text{Gar}^A, T \triangleright G^A} c_T v^T$$

for some $c_T \in \mathcal{O}$.

Proof. In view of (5.4), for each $T \in \text{Gar}^A$, there exists a unique $u^T \in \mathcal{D}^A$ with $T = u^T T^A$. By Corollary 5.12, there exist $d_T \in \mathcal{O}$ such that

$$\begin{aligned} v^{G^A} &= \psi^{G^A} z^\mu = \left(\tau_{u^{G^A}}^A + \sum_{T \in \text{Gar}^A, T \neq G^A} d_T \tau_{u^T}^A \right) z^A \\ &= g^A z^\mu + \sum_{T \in \text{Gar}^A, T \neq G^A} (d_T - 1) \tau_{u^T}^A z^A. \end{aligned}$$

Since $g^A z^\mu = 0$ by Definition 5.9(iv), the result now follows by (inverting the equations in) Corollary 5.12. $\square$

We now make the first step towards describing a standard homogeneous basis of S^μ . In Corollary 6.24 below we show that (5.15) is a basis.

Proposition 5.14. *Let $\mu \in \mathcal{P}_\alpha^\kappa$. The elements of the set*

$$\{v^T \mid T \in \text{St}(\mu)\} \tag{5.15}$$

span S^μ over $\mathcal{O}$. Moreover, we have $\deg(v^T) = \deg(T)$ for all $T \in \text{St}(\mu)$.

Proof. Note that $v^T = \psi^T e(\mathbf{i}^\mu) z^\mu$. Now, using [6, Corollary 3.14], we have $\deg(\psi^T e(\mathbf{i}^\mu)) = \deg(T) - \deg(T^\mu)$, which implies the second statement of the proposition, as $\deg(z^\mu) = \deg(T^\mu)$ by definition.

By Theorem 5.6, it suffices to show that for every row-strict tableau T of shape μ , the vector $v^T \in S^\mu$ is an $\mathcal{O}$ -linear combination of elements in (5.15). We prove this by inverse induction on the Bruhat order on the row-strict tableaux T . The induction starts when $T = T^\mu$, the unique maximal row-strict tableau. In this case T is standard so there is nothing to prove.

For the inductive step, assume that the result has been proved for all row standard tableaux $U \triangleright T$. If the row strict tableau T is standard then there is nothing to prove, so suppose that T is not standard. Then by Lemma 5.2, there exists a Garnir tableaux G of shape μ and $w \in \mathfrak{S}_d$ such that $T = wG$ and $\ell(w^T) = \ell(w^G) + \ell(w)$. Using Proposition 3.15 for the second equality, and then Lemma 5.13 for the last equality, we get

$$\begin{aligned} v^T &= \psi^T e(\mathbf{i}^\mu) z^\mu = (\psi_w \psi^G e(\mathbf{i}^\mu) + x e(\mathbf{i}^\mu)) z^\mu = \psi_w v^G + x z^\mu \\ &= \sum_{T \triangleright G} c_T \psi_w v^T + x z^\mu, \end{aligned}$$

where x is a linear combination of elements of the form $\psi_u f(y)e(\mathbf{i})$ such that $u < w$ and $f(y)$ is a polynomial in $y_1, \dots, y_d$. The result now follows by induction. $\square$

6. CYCLOTOMIC HECKE ALGEBRAS AND SPECHT MODULES

Recall from Definition 5.9 that we have defined by generators and relations the universal graded (row) Specht modules $S^\mu = S^\mu(\mathcal{O})$ for the KLR algebra R_α for all multipartitions $\mu \in \mathcal{P}_\alpha^\kappa$. In this section we connect these universal Specht modules to the usual Specht modules for the affine Hecke algebras H_α via the isomorphism between the cyclotomic quotients of the KLR algebras and of the affine Hecke algebras constructed in [3]. This will allow us to obtain a standard homogeneous basis for $S^\mu(\mathcal{O})$ using [6, 11].

In this section we will need to distinguish between the universal graded (row) Specht modules $S^\mu = S^\mu(\mathcal{O})$ for R_α and the usual graded (row) Specht modules for H_α , which we will denote S_H^μ . The Specht modules S_H^μ are defined as cell modules for the cellular algebra H_α^Λ . We review their properties below.

6.1. Ground field and parameters. Let F be a field, and $\xi \in F^\times$ be an invertible element. Let e be the smallest positive integer such that $1 + \xi + \dots + \xi^{e-1} = 0$, setting $e := 0$ if no such integer exists. This e allows us to use the Lie theoretic notation of section 2.1. In particular, we have $I = \mathbb{Z}/e\mathbb{Z}$, Γ , Q_+ , P_+ , etc.

For $i \in I$ define the scalar $\nu(i) \in F$ as follows:

$$\nu(i) := \begin{cases} i & \text{if } \xi = 1, \\ \xi^i & \text{if } \xi \neq 1. \end{cases} \quad (6.1)$$

6.2. Cyclotomic Hecke algebra. Let $H_d = H_d(F, \xi)$ be the *affine Hecke algebra* over the ground field F associated to the symmetric group $\mathfrak{S}_d$ with parameter ξ . Thus, if $\xi \neq 1$, then H_d is the F -algebra generated by

$$T_1, \dots, T_{d-1}, X_1^{\pm 1}, \dots, X_d^{\pm 1}$$

subject only to the relations

$$T_r^2 = (\xi - 1)T_r + \xi \quad (1 \leq r < d), \quad (6.2)$$

$$T_r T_{r+1} T_r = T_{r+1} T_r T_{r+1} \quad (1 \leq r < d-1), \quad (6.3)$$

$$T_r T_s = T_s T_r \quad (1 \leq r, s < d, |r - s| > 1). \quad (6.4)$$

$$X_r^{\pm 1} X_s^{\pm 1} = X_s^{\pm 1} X_r^{\pm 1} \quad (1 \leq r, s \leq d), \quad (6.5)$$

$$X_r X_r^{-1} = 1 \quad (1 \leq r \leq d), \quad (6.6)$$

$$T_r X_r T_r = \xi X_{r+1} \quad (1 \leq r < d), \quad (6.7)$$

$$T_r X_s = X_s T_r \quad (1 \leq r < d, 1 \leq s \leq d, s \neq r, r+1). \quad (6.8)$$

If $\xi = 1$, then H_d is the F -algebra generated by

$$T_1, \dots, T_{d-1}, X_1, \dots, X_d$$

subject only to the relations (6.2)–(6.4) and the relations:

$$X_r X_s = X_s X_r \quad (1 \leq r, s \leq d), \quad (6.9)$$

$$T_r X_{r+1} = X_r T_r + 1 \quad (1 \leq r < d), \quad (6.10)$$

$$T_r X_s = X_s T_r \quad (1 \leq r < d, 1 \leq s \leq d, s \neq r, r+1). \quad (6.11)$$

Recall that in (2.3) and (2.4) we have fixed a level l , a tuple $\kappa = (k_1, \dots, k_l)$, and the corresponding weight $\Lambda = \Lambda_{k_1} + \dots + \Lambda_{k_l} \in P_+$. The *cyclotomic Hecke algebra* $H_d^\Lambda = H_d^\Lambda(F, \xi)$ is the quotient

$$H_d^\Lambda := H_d / \langle \prod_{i \in I} (X_1 - \nu(i))^{(\Lambda, \alpha_i)} \rangle = H_d / \langle \prod_{m=1}^l (X_1 - \nu(k_m)) \rangle. \quad (6.12)$$

6.3. Weight spaces and idempotents. Let $\mathbf{i} = (i_1, \dots, i_d) \in I^d$, and let M be a finite dimensional H_d^Λ -module. Define the $\mathbf{i}$ -weight space of M as follows:

$$M_{\mathbf{i}} = \{v \in M \mid (X_r - \nu(i_r))^N v = 0 \text{ for } N \gg 0 \text{ and } r = 1, \dots, d\}.$$

It is known (see e.g. [10, Lemma 4.7] and [18, Lemma 7.1.2]) that all eigenvalues of $X_1, \dots, X_d$ in M are of the form $\nu(i)$, for $i \in I$, and so we have a *weight space decomposition*:

$$M = \bigoplus_{\mathbf{i} \in I^d} M_{\mathbf{i}}.$$

Using the weight space decomposition of the left regular H_d^Λ -module, one gets a system of orthogonal idempotents

$$\{e(\mathbf{i}) \mid \mathbf{i} \in I^d\} \quad (6.13)$$

in H_d^Λ , all but finitely many of which are zero, such that $\sum_{\mathbf{i} \in I^d} e(\mathbf{i}) = 1$, and

$$e(\mathbf{i})M = M_{\mathbf{i}} \quad (\mathbf{i} \in I^d)$$

for any finite dimensional H_d^Λ -module M .

If $\alpha \in Q_+$ is of height d , define $e_\alpha := \sum_{\mathbf{i} \in I^\alpha} e(\mathbf{i}) \in H_d^\Lambda$. By [20] and [2, Theorem 1], e_α is either zero or it is a primitive central idempotent in H_d^Λ . Hence the algebra

$$H_\alpha^\Lambda := e_\alpha H_d^\Lambda \quad (6.14)$$

is either zero or it is a single *block* of the algebra H_d^Λ .

6.4. The Isomorphism Theorem. Define elements of H_α^Λ as follows:

$$y_r := \begin{cases} \sum_{\mathbf{i} \in I^\alpha} (1 - \nu(i_r)^{-1} X_r) e(\mathbf{i}), & \text{if } \xi \neq 1, \\ \sum_{\mathbf{i} \in I^\alpha} (X_r - \nu(i_r)) e(\mathbf{i}), & \text{if } \xi = 1, \end{cases} \quad (6.15)$$

for $1 \leq r \leq d$. Next, if $1 \leq r < d$ and $\mathbf{i} \in I^d$ we define

$$\psi_r := \sum_{\mathbf{i} \in I^\alpha} (T_r + P_r(\mathbf{i})) Q_r(\mathbf{i})^{-1} e(\mathbf{i}). \quad (6.16)$$

where $P_r(\mathbf{i})$ and $Q_r(\mathbf{i})^{-1}$ are certain polynomials in $F[y_r, y_{r+1}]$ which are explicitly defined in [3]. This gives us the following elements of H_α^Λ :

$$\{e(\mathbf{i}) \mid \mathbf{i} \in I^\alpha\} \cup \{y_1, \dots, y_d\} \cup \{\psi_1, \dots, \psi_{d-1}\}. \quad (6.17)$$

Note that these elements have the same names as the generators of the KLR algebras in (3.1). This is not a coincidence in view of the following Isomorphism Theorem:

Theorem 6.18. [3, Theorem 1.1] *Suppose that $\alpha \in Q_+$ has height d and $\Lambda \in P_+$. Then H_α^Λ is generated by the elements (6.17) subject only to the relations (3.2)–(3.11). In other words, $H_\alpha^\Lambda(F, \xi) \cong R_\alpha^\Lambda(F)$.*

In what follows we identify $H_\alpha^\Lambda(F, \xi)$ and $R_\alpha^\Lambda(F)$. In particular, $H_\alpha^\Lambda(F, \xi)$ is now $\mathbb{Z}$ -graded.

6.5. Graded Specht modules S_H^μ for Hecke algebras. Let $\alpha \in Q_+$ be of height d and fix a multipartition $\mu \in \mathcal{P}_\alpha^\kappa$. The *graded (row) Specht module* $S_H^\mu = S_H^\mu(F)$ for H_α^Λ is defined in [6]. These graded Specht modules turn out to be the *cell modules* for H_α^Λ considered as a graded cellular algebra as in [11]. We will not need the exact definition, only the following key properties of these modules. Recall the notation of section 2.3.

Lemma 6.19. [14, Proposition 3.7] *Let $\mu \in \mathcal{P}_\alpha^\kappa$. There is a homogeneous generator z_H^μ of S_H^μ with $\deg(z_H^\mu) = \deg(\mathbf{T}^\mu)$, $z_H^\mu \in e(\mathbf{i}^\mu)S_H^\mu$, and $y_r z_H^\mu = 0$, for all $r = 1, \dots, d$.*

Let $\mathbf{T}$ be a μ -tableau. Recall from (3.16) that we have defined the element $\psi^\mathbf{T} = \psi_{w^\mathbf{T}}$ in $R_\alpha^\Lambda(F) = H_\alpha^\Lambda(F)$. Set

$$v_H^\mathbf{T} := \psi^\mathbf{T} z_H^\mu \in S_H^\mu.$$

Just like ψ_w the vector $v_H^\mathbf{T}$ will, in general, depend upon on the choice of preferred reduced decomposition of $w^\mathbf{T} \in \mathfrak{S}_d$. Note that $z_H^\mu = v_H^{\mathbf{T}^\mu}$.

Lemma 6.20. [6, Lemma 4.9] *Let $\mu \in \mathcal{P}_d^\kappa$ and $\mathbf{T} \in \text{St}(\mu)$. If $r \downarrow_{\mathbf{T}} r+1$ or $r \rightarrow_{\mathbf{T}} r+1$ then*

$$\psi_r v_H^\mathbf{T} = \sum_{\mathbf{S} \in \text{St}(\mu), \mathbf{S} \triangleright \mathbf{T}, \mathbf{i}(\mathbf{S}) = \mathbf{i}(\mathbf{s}_r \mathbf{T})} a_{\mathbf{S}} v_H^\mathbf{S}.$$

for some $a_{\mathbf{S}} \in F$. In particular, $\psi_r z_H^\mu = 0$ whenever $r \rightarrow_{\mathbf{T}^\mu} r+1$.

Theorem 6.21. [6] *Suppose that $\mu \in \mathcal{P}_\alpha^\kappa$. Then*

- (i) *For any μ -tableau $\mathbf{T}$ we have $v_H^\mathbf{T} \in e(\mathbf{i}(\mathbf{T}))S_H^\mu$.*
- (ii) *If $\mu \in \text{St}(\mu)$, then $\deg(v_H^\mathbf{T}) = \deg(\mathbf{T})$.*
- (iii) *$\{v_H^\mathbf{T} \mid \mathbf{T} \in \text{St}(\mu)\}$ is a basis of $S_H^\mu(F)$. Moreover, for any μ -tableau $\mathbf{T}$, we have*

$$v_H^\mathbf{T} = \sum_{\substack{\mathbf{S} \in \text{St}(\mu) \\ \mathbf{S} \triangleright \mathbf{T}, \mathbf{i}(\mathbf{S}) = \mathbf{i}(\mathbf{T})}} b_{\mathbf{S}} v_H^\mathbf{S}$$

for some constants $b_{\mathbf{S}} \in F$.

The following corollary should be compared with Lemma 5.13.

Corollary 6.22. *Suppose that $\mu \in \mathcal{P}_d^\kappa$ and that $\mathbf{G} = \mathbf{G}^A$, where $A \in \mu$ is a Garnir node. Then*

$$v_H^\mathbf{G} = \sum_{\mathbf{T} \in \text{Gar}^A, \mathbf{T} \triangleright \mathbf{G}} c_{\mathbf{T}} v_H^\mathbf{T}$$

for some $c_{\mathbf{T}} \in F$.

Proof. This comes from Theorem 6.21 and Lemma 5.5. □

6.6. Connecting the universal row Specht modules with the cell modules. Since we have identified the algebras $H_\alpha^\Lambda(F)$ and $R_\alpha^\Lambda(F)$ we may consider the Specht modules $S_H^\mu(F)$ as an $R_\alpha^\Lambda(F)$ -module. Inflating from $R_\alpha^\Lambda(F)$ to the affine KLR algebra $R_\alpha(F)$, we can now consider $S_H^\mu(F)$ as an $R_\alpha(F)$ -module. The following theorem shows that, as a graded $R_\alpha(F)$ -module, $S_H^\mu(F)$ is isomorphic to the universal row Specht module $S^\mu(F)$ from Definition 5.9.

Theorem 6.23. *Let $\mu \in \mathcal{P}_\alpha^\kappa$. Then the linear map, which sends the basis elements $v_H^\mathbf{T} \in S_H^\mu(F)$ to $v^\mathbf{T} \in S^\mu(F)$ for all $\mathbf{T} \in \text{St}(\mu)$, is a homogeneous isomorphism $S_H^\mu(F) \xrightarrow{\sim} S^\mu(F)$ of graded $R_\alpha(F)$ -modules.*

Proof. In this proof all modules and algebras are vector spaces over F , so we will suppress F from our notation. We will construct the isomorphism in the other direction: $S^\mu \xrightarrow{\sim} S_H^\mu$. By Lemmas 6.19, 6.20 and the defining relations for M^μ , cf. (3.22), there exists a surjective degree zero homogeneous homomorphism $\pi : M^\mu \twoheadrightarrow S_H^\mu$ of graded R_α -modules which maps $m^\mathbf{T}$ to $v_H^\mathbf{T}$ for any row-strict μ -tableau $\mathbf{T}$. By Theorem 6.21 and Proposition 5.14, it now suffices to check that the homogeneous Garnir relations $g^A z_H^\mu = 0$ hold in S_H^μ , for all Garnir nodes $A \in \mu$.

Fix a Garnir node A . Let $k = k^A$, $f = f^A$ and $\mathfrak{S}^A \cong \mathfrak{S}_k$ be the brick permutation group defined in section 5.2. By Corollary 5.12, $\{m^\mathbf{T} \mid \mathbf{T} \in \text{Gar}^A\}$ is an F -basis of $T^{\mu,A}$. Note that $\mathbf{G}^A$ is the only non-standard tableaux in Gar^A . As $\pi(T^{\mu,A})$ is spanned by the vectors $\{v_H^\mathbf{T} = \pi(m^\mathbf{T}) \mid \mathbf{T} \in \text{Gar}^A\}$, Corollary 6.22 shows that $\{v_H^\mathbf{T} \mid \mathbf{T} \in \text{Gar}^A \setminus \{\mathbf{G}^A\}\}$ is a basis of $\pi(T^{\mu,A})$. So $\dim \pi(T^{\mu,A}) = \dim T^{\mu,A} - 1$.

Recall from Theorem 5.11 that the group $\mathfrak{S}^A$ acts on the brick permutation subspace $T^{\mu,A}$ with its simple reflections acting as $\tau_1^A, \dots, \tau_{k-1}^A$. Moreover, with respect to this action, $T^{\mu,A} \cong \text{ind}_{F\mathfrak{S}_{(f,k-f)}}^{F\mathfrak{S}_k} F_{(f,k-f)}$. Since the elements of $\mathfrak{S}^A$ act on $T^{\mu,A}$ as specific elements of R_α , and π is an R_α -homomorphism, π induces an $F\mathfrak{S}^A$ -homomorphism $T^{\mu,A} \rightarrow \pi(T^{\mu,A})$. By the dimension observations in the previous paragraph, the kernel of this map is a one dimensional $F\mathfrak{S}_k$ submodule of $T^{\mu,A} = \text{ind}_{F\mathfrak{S}_{(f,k-f)}}^{F\mathfrak{S}_k} F_{(f,k-f)}$. Therefore, unless $k = 2$, $f = 1$, and $\text{char } F \neq 2$, this kernel is the unique trivial submodule of $\text{ind}_{F\mathfrak{S}_{(f,k-f)}}^{F\mathfrak{S}_k} F_{(f,k-f)}$. Hence, in this case, $\ker \pi$ is spanned by $\sum_{u \in \mathcal{Q}^A} \tau_u^A m^A = g^A m^\mu$. Hence $g^A z_H^\mu = \pi(g^A m^\mu) = 0$, so that the Garnir relation holds in the Specht module S_H^μ , as desired.

It remains to consider the exceptional case $k = 2$, $f = 1$, $\text{char } F \neq 2$. In this case we claim that $(\tau_1^A + 1)z^A = 0$, for this we need to rule out the possibility that $(\tau_1^A - 1)z^A = 0$. Since $\tau_1^A z^A = (\sigma_1^A + 1)z^A$, we just need to prove that $\sigma_1^A z^A \neq 0$. Let $A = (a, b, m)$, and r be the entry which occupies the node $(a + 1, b, m)$ in $\mathbf{G}^A$. But by Lemma 4.5, we have

$$\psi_r \sigma_1^A z^A = -2\psi_r z^A = -2v^{s_r \mathbf{T}^A} \neq 0,$$

since the tableau $s_r \mathbf{T}^A$ is standard and $\text{char } F \neq 2$. $\square$

We can now improve on Proposition 5.14:

Corollary 6.24. *Let $\mu \in \mathcal{P}_\alpha^\kappa$. Then the universal row Specht module $S^\mu(\mathcal{O})$ for $R_\alpha(\mathcal{O})$ has $\mathcal{O}$ -basis*

$$\{v^\mathbf{T} \mid \mathbf{T} \in \text{St}(\mu)\}. \quad (6.25)$$

Proof. As $S^\mu(\mathcal{O}) \cong S^\mu(\mathbb{Z}) \otimes_{\mathbb{Z}} \mathcal{O}$, we may assume that $\mathcal{O} = \mathbb{Z}$. By Proposition 5.14, the elements (6.25) span $S^\mu(\mathbb{Z})$. Suppose that we have a relation $\sum_{\mathbf{T} \in \text{St}(\mu)} c_\mathbf{T} v^\mathbf{T} = 0$ with $c_\mathbf{T} \in \mathbb{Z}$. Extending scalars to $\mathbb{C}$, we get the relation $\sum_{\mathbf{T} \in \text{St}(\mu)} c_\mathbf{T} v^\mathbf{T} = 0$ in $S^\mu(\mathbb{C})$. Pick a parameter $\xi \in \mathbb{C}$ which is a primitive e th root of unity in $\mathbb{C}$ if $e > 0$ and not a root of unity if $e = 0$. Then by Theorem 6.23, we get the relation $\sum_{\mathbf{T} \in \text{St}(\mu)} c_\mathbf{T} v_H^\mathbf{T} = 0$ in $S_H^\mu(\mathbb{C})$, which is

the usual Specht module for $H_\alpha^\Lambda(\mathbb{C}, \xi)$. By Theorem 6.21(iii), $c_{\mathbf{T}} = 0$ for all $\mathbf{T} \in \text{St}(\mu)$. $\square$

Corollary 6.26. *Let $\mu \in \mathcal{P}_\alpha^\kappa$. The universal row Specht module $S^\mu(\mathcal{O})$ factors through the natural surjection $R_\alpha(\mathcal{O}) \twoheadrightarrow R_\alpha^\Lambda(\mathcal{O})$ so that $S^\mu(\mathcal{O})$ is naturally a graded $R_\alpha^\Lambda(\mathcal{O})$ -module.*

Proof. In view of (3.11), we just need to prove that

$$y_1^{(\Lambda, \alpha_{i_1})} e(\mathbf{i}) S^\mu(\mathcal{O}) = 0 \quad (\mathbf{i} = (i_1, \dots, i_d) \in I^\alpha).$$

We may assume that $\mathcal{O} = \mathbb{Z}$. Next, since $S^\mu(\mathbb{Z}) \hookrightarrow S^\mu(\mathbb{Z}) \otimes \mathbb{C} = S^\mu(\mathbb{C})$, we may now assume that $\mathcal{O} = \mathbb{C}$. Choose $\xi \in \mathbb{C}$ as in the proof of Corollary 6.24. Then by Theorem 6.23, we have $S^\mu(\mathbb{C}) = S_H^\mu(\mathbb{C})$, which is the usual Specht module for $H_\alpha^\Lambda(\mathbb{C}, \xi)$. Hence, $S^\mu(\mathbb{C})$ is a $R_\alpha^\Lambda(\mathbb{C})$ module since $H_\alpha^\Lambda(\mathbb{C}, \xi) \cong R_\alpha^\Lambda(\mathbb{C})$. Hence, the action of $R_\alpha^\Lambda(\mathbb{C})$ satisfies the cyclotomic relation (3.11), implying that $S^\mu(\mathcal{O})$ is an $R_\alpha^\Lambda(\mathcal{O})$ -module as we wanted to show. $\square$

Now the following is clear:

Corollary 6.27. *Let $\mu \in \mathcal{P}_\alpha^\kappa$.*

- (i) *As a graded $R_\alpha^\Lambda(\mathcal{O})$ -module, the universal row Specht module $S^\mu(\mathcal{O})$ is generated by the homogeneous element z^μ of degree $\deg(\mathbf{T}^\mu)$ subject only to the relations (i)–(iv) from Definition 5.9.*
- (ii) *As a graded H_α^Λ -module, the row Specht module S_H^μ is generated by the homogeneous element z^μ of degree $\deg(\mathbf{T}^\mu)$ subject only to the relations (i)–(iv) from Definition 5.9.*

7. COLUMN SPECHT MODULES

Having a presentation for a module does not automatically imply a presentation for the dual module. In this section, we define a column version S_μ of the universal graded Specht module corresponding to a multipartition μ . Then in Theorem 7.25 we show that the universal column Specht module S_μ is isomorphic to (a degree shift of) the homogeneous dual $(S^\mu)^\circledast$ of the universal row Specht module S^μ .

In the section we again work over an arbitrary commutative unital ground ring $\mathcal{O}$, unless otherwise stated. We fix $\alpha \in Q_+$, $\mu \in \mathcal{P}_\alpha^\kappa$, and set $d := \text{ht}(\alpha)$.

7.1. Column block intertwiners. In this section we assume that $e > 0$. Recall from (2.2) that δ is the null root and observe that $\delta' = \delta$ in the notation of section 8.2. Therefore, sgn is an automorphism of $R_{k\delta}$, see (3.14).

Fix $i \in I$ and a composition $\lambda = (\lambda_1, \dots, \lambda_n)$ of k . Define

$$\vec{\mathbf{s}}(i, -\lambda) := (\mathbf{s}(i, -e\lambda_1), \dots, \mathbf{s}(i, -e\lambda_n)).$$

We consider the corresponding permutation module $M(i, -\lambda) := M(\vec{\mathbf{s}}(i, -\lambda))$ for $R_{k\delta}$ as in section 3.6. Let $\mathbf{j} = (j_1, \dots, j_{ke}) := \mathbf{j}(\vec{\mathbf{s}}(i, -\lambda))$ as defined in (3.21). We have $\mathbf{j} = \mathbf{s}(i, -ke)$. Let $e(i, -\lambda) := e(\mathbf{j})$ and $m(i, -\lambda) = m(\vec{\mathbf{s}}(i, -\lambda)) \in M(i, -\lambda)$ as in (3.20).

Recall from section 8.2 that if M is an $R_{k\delta}$ -module then M^{sgn} is the $R_{k\delta}$ -module obtained from M by twisting with the sign automorphism sgn .

Lemma 7.1. *We have*

- (i) $\text{sgn}(e(i, -\lambda)) = e(-i, \lambda)$.

- (ii) *There is an isomorphism $M(i, -\lambda) \cong M(-i, \lambda)^{\text{sgn}}$ of graded $R_{k\delta}$ -modules, under which $m(i, -\lambda)$ corresponds to $m(-i, \lambda)$.*

Proof. (i) is clear from (3.14), and (ii) is clear from (3.22). $\square$

From section 4, we have the elements $w_r \in \mathfrak{S}_{ke}$, $\sigma_r = \psi_{w_r}e(-i, \lambda)$ and $\tau_r = (\sigma_r + 1)e(-i, \lambda)$, for $1 \leq r < k$. Set

$$\sigma^r := (-1)^e \psi_{w_r}e(i, -\lambda) \quad \text{and} \quad \tau^r = (\sigma^r + 1)e(i, -\lambda),$$

for $r = 1, \dots, k-1$, and define the *column block permutation subspace* $T(i, -\lambda) \subseteq M(i, -\lambda)$ to be the $\mathcal{O}$ -span of all vectors of the form

$$\sigma^{r_1} \dots \sigma^{r_a} m(i, -\lambda). \quad (7.2)$$

Lemma 7.3. *We have*

- (i) $\text{sgn}(\sigma^r) = \sigma_r$ and $\text{sgn}(\tau^r) = \tau_r$, for $1 \leq r < k$.
- (ii) *Under the isomorphism of Lemma 7.1(ii), $T(i, -\lambda)$ corresponds to $T(-i, \lambda)$*

Proof. Since $\ell(w_r) = e^2$, we have $\text{sgn}(\psi_{w_r}) = (-1)^{e^2} \psi_{w_r} = (-1)^e \psi_{w_r}$. So Lemma 7.1(i) yields (i). Part (ii) follows from (i) and Lemma 7.1(ii). $\square$

Lemma 7.3 and Theorem 4.12 now imply the following.

Proposition 7.4. *Suppose that $1 \leq r, s < k$ and $v \in T(i, -\lambda)$. Then*

- (i) $(\tau^r)^2 v = v$.
- (ii) *If $|r - s| > 1$ then $\tau^r \tau^s v = \tau^s \tau^r v$.*
- (iii) *If $r < k-1$ then $\tau^r \tau^{r+1} \tau^r v = \tau^{r+1} \tau^r \tau^{r+1} v$.*

Consequently, $\mathfrak{S}_k$ acts on $T(i, -\lambda)$, and the elements $\tau^u m(i, -\lambda)$ for $u \in \mathfrak{S}_k$ are well-defined. Finally, $T(i, -\lambda) \cong \text{ind}_{\mathcal{O}_{\mathfrak{S}_\lambda}}^{\mathcal{O}_{\mathfrak{S}_k}} \mathcal{O}_\lambda$ as $\mathcal{O}_{\mathfrak{S}_k}$ -modules, and $T(i, -\lambda)$ has $\mathcal{O}$ -basis $\{\tau^u m(i, -\lambda) \mid u \in \mathcal{D}_\lambda\}$.

7.2. Column Garnir tableau. We now rework the combinatorics of row Garnir tableaux for column Garnir tableaux. A node $A = (a, b, m) \in \mu$ is a *column Garnir node* of μ if $(a, b+1, m)$ is a node of μ . The (column) *A-Garnir belt* $\mathbf{B}_A$ is the set of nodes

$$\mathbf{B}_A = \{(c, b, m) \in \mu \mid c \geq a\} \cup \{(c, b+1, m) \in \mu \mid c \leq a\}.$$

Recall from (2.16) that if $\mathbf{T} \in \text{St}(\mu)$ then $\mathbf{T}_\mu \trianglelefteq \mathbf{T}$ and $w_{\mathbf{T}} \in \mathfrak{S}_d$ is the permutation such that $\mathbf{T} = w_{\mathbf{T}} \mathbf{T}_\mu$. Let $u = \mathbf{T}_\mu(a, b, m)$ and $v = \mathbf{T}_\mu(a+1, b, m)$. The (column) *A-Garnir tableau* $\mathbf{G}_A$ is the μ -tableaux which agrees with $\mathbf{T}_\mu$ outside of $\mathbf{B}_A$ and where the numbers $u, u+1, \dots, v$ are inserted into the Garnir belt in order, from top right to left bottom.

Just as in section 5.1 we have the following two results.

Lemma 7.5. *Suppose that $A \in \mu$ is a column Garnir node and $\mathbf{S} \in \text{St}(\mu)$. If $\mathbf{G}_A \triangleleft \mathbf{S}$ then $\mathbf{S}$ agrees with $\mathbf{T}_\mu$ outside of $\mathbf{B}_A$.*

Lemma 7.6. *Suppose that $\mathbf{T}$ is a column strict μ -tableaux which is not standard. Then there exists a column Garnir tableaux $\mathbf{G}$ and $w \in S_d$ such that $\mathbf{T} = w\mathbf{G}$ and $\ell(w_{\mathbf{T}}) = \ell(w_{\mathbf{G}}) + \ell(w)$.*

7.3. Column bricks. A (column A -)brick is a set of e nodes

$$\{(c, d, m), (c+1, d, m), \dots, (c+e-1, d, m)\} \subseteq \mathbf{B}_A$$

such that $\text{res}(c, d, m) = \text{res } A$. The Garnir belt $\mathbf{B}_A$ is a disjoint union of the bricks that it contains together with less than e nodes at the bottom of column a which are not contained in a brick and less than e nodes at the top of column $a+1$ which are not contained in a brick.

For example, if $e = 2$, then the $(3, 1, 2)$ -Garnir belt of $\mu = ((1), (7, 7, 4, 1))$ contains two bricks:

$$\mathbb{G}_A = \begin{array}{|c|c|c|c|c|c|c|} \hline & & & & & & \\ \hline & 1 & 3 & 8 & 11 & 14 & 16 & 18 \\ \hline & 2 & 4 & 9 & 12 & 15 & 17 & 19 \\ \hline 6 & 5 & 10 & 13 & & & & \\ \hline 7 & & & & & & & \\ \hline \end{array}.$$

Let $k = k_A$ be the number of bricks in $\mathbf{B}_A$. Label the bricks $B_A^1, B_A^2, \dots, B_A^k$ in $\mathbf{B}_A$ from top to bottom first down column $b+1$ and then down column b of μ . Set $k = 0$ if $\mathbf{B}_A$ does not contain any bricks.

If $k > 0$ let $n = n_A$ be the smallest number in $\mathbb{G}_A$ which is contained in a brick in $\mathbf{B}_A$. In the example above, $k = 2$ and $n = 4$. Define

$$w_A^r = \prod_{a=n+re-e}^{n+re-1} (a, a+e) \in \mathfrak{S}_d \quad (1 \leq r < k).$$

The (column) brick permutation group is the subgroup $\mathfrak{S}_A$ of $\mathfrak{S}_d$ generated by $w_A^1, w_A^2, \dots, w_A^{k-1}$. Then $\mathfrak{S}_A \cong \mathfrak{S}_k$.

Let Gar_A be the set of all column-strict μ -tableaux which are obtained from the Garnir tableau $\mathbb{G}_A$ by acting with the brick permutation group $\mathfrak{S}_A$ on $\mathbb{G}_A$. All tableaux in Gar_A are standard except for $\mathbb{G}_A$, $\mathbb{G}_A$ is the maximal element of Gar_A , and there is a unique minimal tableau $\mathbf{T}_A$ in Gar_A . If $\mathbf{T} \in \text{Gar}_A$ then $\mathbf{i}(\mathbf{T}) = \mathbf{i}(\mathbb{G}_A)$. We let $\mathbf{i}_A := \mathbf{i}(\mathbb{G}_A)$.

Define $f = f_A$ to be the number of A -bricks in column b of the Garnir belt $\mathbf{B}_A$ and let $\mathcal{D}_A$ be the set of minimal length left coset representations of $\mathfrak{S}_f \times \mathfrak{S}_{k-f}$ in $\mathfrak{S}_A \cong \mathfrak{S}_k$. Just as in (5.4), we have

$$\text{Gar}_A = \{w\mathbf{T}_A \mid w \in \mathcal{D}_A\}. \quad (7.7)$$

Finally, as in Lemma 5.5, we have:

Lemma 7.8. *Let $A \in \mu$ be a column Garnir node. Then*

$$\text{Gar}_A \setminus \{\mathbb{G}_A\} = \{\mathbf{T} \in \text{St}(\mu) \mid \mathbf{T} \trianglelefteq \mathbb{G}_A \text{ and } \mathbf{i}(\mathbf{T}) = \mathbf{i}_A\}.$$

Moreover, $\text{codeg}(\mathbf{T}) = \text{codeg}(\mathbb{G}_A)$ for all $\mathbf{T} \in \text{Gar}_A$.

7.4. The column permutation modules M_μ . Let $C_1, \dots, C_g$ be the non-empty columns of μ counted from left to right in the component $\mu^{(l)}$, then from left to right in the component $\mu^{(l-1)}$, and so on, until from left to right in the component $\mu^{(1)}$ of μ . We emphasize that the order of the components of μ is reversed here.

To each $1 \leq a \leq g$ we associate the segment $\mathbf{c}(a) := \mathbf{s}(i, -N)$, where the column C_a has length N and i is the residue of the top node of C_a . Let $\vec{\mathbf{c}} = (\mathbf{c}(1), \dots, \mathbf{c}(g))$, and, recalling the definitions from section 3.6, set

$$M_\mu = M_\mu(\mathcal{O}) := M(\vec{\mathbf{c}}) \langle \text{codeg } \mathbf{T}_\mu \rangle.$$

The module M_μ is generated by the vector $m_\mu := m(\vec{\mathbf{c}})$ of degree $\text{codeg } \mathbf{T}_\mu$. For any μ -tableau $\mathbf{T}$, define $m_{\mathbf{T}} := \psi_{\mathbf{T}} m_\mu$. As a special case of Theorem 3.23, we have:

Theorem 7.9. $\{m_{\mathbf{T}} \mid \mathbf{T} \text{ is a column-strict } \mu\text{-tableau}\}$ is an $\mathcal{O}$ -basis of M_μ .

7.5. Universal column Specht modules S_μ . Fix a column Garnir node $A \in \mu$, and let $\mathfrak{S}_A = \langle w_A^1, \dots, w_A^{k-1} \rangle$ be the corresponding block permutation group. For any $\mathbf{S} \in \text{Gar}_A$, we can write $w_{\mathbf{S}} = u_{\mathbf{S}} w_{\mathbf{T}_A}$ with $\ell(w_{\mathbf{S}}) = \ell(u_{\mathbf{S}}) + \ell(w_{\mathbf{T}_A})$ and $u_{\mathbf{S}} \in \mathcal{D}_A$. By Lemma 3.17, $w_{\mathbf{S}}$, $u_{\mathbf{S}}$, and $w_{\mathbf{T}_A}$ are fully commutative so we have elements $\psi_{\mathbf{S}}$, $\psi_{u_{\mathbf{S}}}$ and $\psi_{\mathbf{T}_A}$, with $\psi_{\mathbf{S}} = \psi_{u_{\mathbf{S}}} \psi_{\mathbf{T}_A}$, each of which is independent of the choice of preferred decomposition.

Set $m_A := m_{\mathbf{T}_A} = \psi_{\mathbf{T}_A} m_\mu$ and define

$$\sigma_A^r := (-1)^e \psi_{w_A^r} e(\mathbf{i}_A) \quad \text{and} \quad \tau_A^r := (\sigma_A^r + 1) e(\mathbf{i}_A).$$

Any element $u \in \mathfrak{S}_A$ can be written as a reduced product $u = w_{r_1}^A \dots w_{r_m}^A$. If u is fully commutative then $\tau_A^u := \tau_A^{r_1} \dots \tau_A^{r_m}$ is independent of the choice of the reduced expression by Lemma 3.17, so we have well-defined elements $\{\tau_A^u \mid u \in \mathcal{D}_A\}$.

Definition 7.10. Suppose that $A \in \mu$ is a column Garnir node. The *column Garnir element* is

$$g_A := \sum_{u \in \mathcal{D}_A} \tau_A^u \psi_{\mathbf{T}_A} \in R_\alpha.$$

Since $\psi_{\mathbf{T}_A} m_\mu = m_A$, we have $g_A m_\mu = \sum_{u \in \mathcal{D}_A} \tau_A^u m_A$, and, by Lemma 7.8, all summands on the right hand side have the same degree. If $k = 0$ then $\mathcal{D}_A = \{1\}$, $\mathbf{G}_A = \mathbf{T}_A$ and $g_A = \psi_{\mathbf{G}_A}$.

Definition 7.11. The *universal graded column Specht module* $S_\mu = S_\mu(\mathcal{O})$ is the graded R_α -module generated by the vector z_μ of degree $\text{codeg}(\mathbf{T}_\mu)$ subject only to the following relations:

- (i) $e(\mathbf{j}) z_\mu = \delta_{\mathbf{j}, \mathbf{i}_\mu} z_\mu$ for all $\mathbf{j} \in I^\alpha$;
- (ii) $y_r z_\mu = 0$ for all $r = 1, \dots, d$;
- (iii) $\psi_r z_\mu = 0$ for all $r = 1, \dots, d-1$ such that $r \downarrow_{\mathbf{T}_\mu} r+1$;
- (iv) (*homogeneous (column) Garnir relations*) $g_A z_\mu = 0$ for all (column) Garnir nodes A in μ .

In other words, $S_\mu = (R_\alpha / J_{\alpha, \mu}) \langle \text{codeg}(\mathbf{T}_\mu) \rangle$, where $J_{\alpha, \mu}$ is the left ideal of R_α generated by the elements

- (i) $e(\mathbf{j}) - \delta_{\mathbf{j}, \mathbf{i}_\mu}$ for all $\mathbf{j} \in I^\alpha$;
- (ii) y_r for all $r = 1, \dots, d$;
- (iii) ψ_r for all $r = 1, \dots, d-1$ such that $r \downarrow_{\mathbf{T}_\mu} r+1$;
- (iv) g_A for all column Garnir nodes $A \in \mu$.

Since the elements (i)-(iii) generate the left ideal K_μ with $R_\alpha / K_\mu \cong M_\mu$, we have a natural surjection $M_\mu \twoheadrightarrow S_\mu$, which maps m_μ to z_μ , and the kernel $J_\mu = J_{\alpha, \mu} m_\mu$ of this surjection is generated by the Garnir relations.

7.6. Column brick permutation space $T_{\mu,A}$. The (column) brick permutation space $T_{\mu,A} \subseteq M_\mu$ is the $\mathcal{O}$ -span of all elements of the form $\sigma_A^{r_1} \dots \sigma_A^{r_a} m_A$. Repeating the argument of Theorem 5.11 now gives:

Theorem 7.12. *Suppose that $A \in \mu$ is a column Garnir node, and let $k = k_A$ and $f = f_A$. Then:*

- (i) $T_{\mu,A}$ the $\mathcal{O}$ -span of all elements of the form $\tau_A^{r_1} \dots \tau_A^{r_a} m_A$. In particular, the elements $\tau_A^1, \dots, \tau_A^{k-1}$ act on $T_{\mu,A}$.
- (ii) As $\mathcal{O}$ -linear operators on $T_{\mu,A}$, the elements $\tau_A^1, \dots, \tau_A^{k-1}$ satisfy the Coxeter relations for the symmetric group $\mathfrak{S}_k$. Thus, we can consider $T_{\mu,A}$ as an $\mathcal{O}\mathfrak{S}_k$ -module.
- (iii) There is an isomorphism of $\mathcal{O}\mathfrak{S}_k$ -modules

$$T_{\mu,A} \cong \text{ind}_{\mathcal{O}\mathfrak{S}_{(f,k-f)}}^{\mathcal{O}\mathfrak{S}_k} \mathcal{O}_{(f,k-f)}$$

under which m_A corresponds to the natural cyclic generator of the induced module on the right hand side.

- (iv) $\{\tau_A^u m_A \mid u \in \mathcal{D}_A\}$ is an $\mathcal{O}$ -basis of $T_{\mu,A}$.

Corollary 7.13. *Suppose that $A \in \mu$ is a column Garnir node of μ , and $S = uT_A \in \text{Gar}_A$ for some $u \in \mathcal{D}_A$. Then*

$$\psi_S m_\mu = \tau_A^u m_A + \sum_{w \in \mathcal{D}_A, w \triangleleft u} c_w \tau_A^w m_A$$

for some $c_w \in \mathcal{O}$. In particular, $\{m_T \mid T \in \text{Gar}_A\}$ is an $\mathcal{O}$ -basis of $T_{\mu,A}$.

7.7. A spanning set for the universal column Specht module. Recall from section 7.5 that $S_\mu \cong M_\mu/J_\mu$ and $z_\mu = m_\mu + J_\mu$. Also set $z_A := m_A + J_\mu \in S_\mu$ for any column Garnir node $A \in \mu$. Recall from (3.16) that for each μ -tableau T we have defined the element $\psi_T \in R_\alpha$, which depends on a fixed choice of reduced decomposition of $w_T \in \mathfrak{S}_d$. We associate to T the homogeneous element $v_T := \psi_T z_\mu \in S_\mu$.

Adapting the arguments from section 5.6 we obtain the following result.

Proposition 7.14. *The elements $\{v_T \mid T \in \text{St}(\mu)\}$ span S_μ over $\mathcal{O}$. Moreover, we have $\deg(v_T) = \text{codeg}(T)$ for all $T \in \text{St}(\mu)$.*

7.8. Graded column Specht modules S_μ^H for Hecke algebras. The graded column Specht modules for the cyclotomic Hecke algebra H_α^Λ were defined in [11, §6] as cell modules for certain graded cellular structure on H_α^Λ (different from the one used to define cell modules S_H^μ). We review the key properties of these modules, paralleling section 6.5.

Recall the definition of the conjugate multipartition μ' and conjugate tableaux from section 2.3. If $\mu \in \mathcal{P}_\alpha^\kappa$ then in general $\mu' \notin \mathcal{P}_\alpha^\kappa$. Because of this we will use a *different labelling of the column Specht modules than* [11]. Let $S_\mu^{HM}(F)$ be the graded column Specht module for $H_\alpha^\Lambda(F)$ constructed in [11, §6.4]. That is, $S_\mu^{HM}(F)$ is a graded cell module with basis $\{\psi'_T \mid T \in \text{St}(\mu)\}$, where $\deg(\psi'_T) = \text{codeg}(T')$ for $T \in \text{St}(\mu)$, using the cellular basis notation of [11, §6.4]. Define

$$S_\mu^H(F) := S_{\mu'}^{HM}(F) \quad \text{and} \quad z_\mu^H := \psi'_{T_{\mu'}}.$$

The following lemma was proved in [14, Proposition 3.7]:

Lemma 7.15. *Let $\alpha \in Q_+$, $d = \text{ht}(\alpha)$, and $\mu \in \mathcal{P}_\alpha^\kappa$. As an H_α^Λ -module, S_μ^H is generated by z_μ^H , $\deg(z_\mu^H) = \text{codeg}(\mathbf{T}_\mu)$, $z_\mu^H \in e(\mathbf{i}_\mu)S_\mu^H$, and $y_r z_\mu^H = 0$, for all $r = 1, \dots, d$.*

Proof. Since $(\mathbf{T}^{\mu'})' = \mathbf{T}_\mu$, the first three claims follow from [11, Proposition 6.10]. That $y_r z_\mu^H = 0$ for all r , can be deduced from [11, (6.2)] and (6.15). Alternatively, it is a special case of [12, Corollary 3.11]. $\square$

For each μ -tableau $\mathbf{T}$, define $v_{\mathbf{T}}^H = \psi_{\mathbf{T}} z_\mu^H \in S_\mu^H$. By [11, Definition 6.9], $v_{\mathbf{T}}^H$ is the same as the element $\psi'_{\mathbf{T}}$ in the notation of [11]. In particular, $z_\mu^H = v_{\mathbf{T}_\mu}^H$.

Lemma 7.16. *Suppose that $\mathbf{T} \in \text{St}(\mu)$. If $r \downarrow_{\mathbf{T}} r+1$ or $r \rightarrow_{\mathbf{T}} r+1$ then*

$$\psi_r v_{\mathbf{T}}^H = \sum_{\mathbf{S} \in \text{St}(\mu), \mathbf{S} \triangleleft \mathbf{T}, \mathbf{i}(\mathbf{S}) = \mathbf{i}(s_r \mathbf{T})} a_{\mathbf{S}} v_{\mathbf{S}}^H.$$

for some $a_{\mathbf{S}} \in F$. In particular, $\psi_r z_\mu^H = 0$ whenever $r \downarrow_{\mathbf{T}_\mu} r+1$.

Proof. This can be deduced from [11, Proposition 6.10(c)] using standard properties of the (ungraded) dual Murphy basis. Alternatively, it follows immediately from [12, Corollary 3.12]. $\square$

The next result is the analogue of Theorem 6.21.

Theorem 7.17. *We have*

- (i) *If $\mathbf{T}$ is a μ -tableau then $v_{\mathbf{T}}^H \in e(\mathbf{i}(\mathbf{T}))S_\mu^H$.*
- (ii) *If $\mathbf{T} \in \text{St}(\mu)$ then $\deg(v_{\mathbf{T}}^H) = \text{codeg}(\mathbf{T})$.*
- (iii) *$\{v_{\mathbf{T}}^H \mid \mathbf{T} \in \text{St}(\mu)\}$ is a basis of $S_\mu^H(F)$. Moreover, for any μ -tableau $\mathbf{T}$,*

$$v_{\mathbf{T}}^H = \sum_{\mathbf{S} \in \text{St}(\mu), \mathbf{S} \trianglelefteq \mathbf{T}, \mathbf{i}(\mathbf{S}) = \mathbf{i}(\mathbf{T})} b_{\mathbf{S}} v_{\mathbf{S}}^H$$

for some constants $b_{\mathbf{S}} \in F$.

Proof. Everything except for the second part of (iii) is clear from results in [11] and the remarks above. Part (iii) can be deduced from [11, Proposition 6.10(c)]. Alternatively, it can be deduced from [12, Theorem 3.9]. $\square$

Corollary 7.18. *Suppose that $A \in \mu$ is a column Garnir node. Then*

$$v_{\mathbf{G}_A}^H = \sum_{\mathbf{T} \in \text{Gar}^A, \mathbf{T} \triangleleft \mathbf{G}_A} c_{\mathbf{T}} v_{\mathbf{T}}^H$$

for some $c_{\mathbf{T}} \in F$.

7.9. Connecting the universal column Specht modules with the cell modules. As in the last section let $S_\mu^H(F)$ be the graded column Specht module for H_α^Λ , where F is a field. As in section 6.6 we consider $S_\mu^H(F)$ as an $R_\alpha(F)$ -module.

Mimicking the proof of Theorem 6.23 and using, in particular, the results in section 7.8, Theorem 7.12 and Corollary 7.18, we can now show that $S_\mu^H(F) \cong S_\mu(F)$ as an $R_\alpha(F)$ -module. As the argument is similar to the proof of Theorem 6.23 we leave the details to the reader.

Theorem 7.19. *There is a homogeneous isomorphism $S_\mu^H(F) \xrightarrow{\sim} S_\mu(F)$ of graded $R_\alpha(F)$ -modules, which maps $v_T^H \in S_\mu^H(F)$ to $v_T \in S_\mu(F)$ for all $T \in \text{St}(\mu)$.*

The following three Corollaries of Theorem 7.19 are proved in exactly the same way as the corresponding results in section 6.6.

Corollary 7.20. *$\{v_T \mid T \in \text{St}(\mu)\}$ is an $\mathcal{O}$ -basis of $S_\mu(\mathcal{O})$.*

Corollary 7.21. *The universal column Specht module $S_\mu(\mathcal{O})$ factors through the natural surjection $R_\alpha(\mathcal{O}) \rightarrow R_\alpha^\Lambda(\mathcal{O})$ so that $S_\mu(\mathcal{O})$ is naturally a graded $R_\alpha^\Lambda(\mathcal{O})$ -module.*

Corollary 7.22. *We have*

- (i) *As a graded $R_\alpha^\Lambda(\mathcal{O})$ -module, the universal column Specht module $S_\mu(\mathcal{O})$ is generated by the homogeneous element z_μ of degree $\text{codeg}(\mathbf{T}_\mu)$ subject only to the relations (i)–(iv) from Definition 7.11.*
- (ii) *As a graded H_α^Λ -module, the column Specht module S_μ^H is generated by the homogeneous element z_μ of degree $\text{codeg}(\mathbf{T}_\mu)$ subject only to the relations (i)–(iv) from Definition 7.11.*

7.10. Contragredient duality for Specht modules. Recall from section 3.2 that $M^\circledast$ denotes the graded dual of the R_α -module M . We now use [11] to show that $S^\mu(\mathcal{O})^\circledast \cong S_\mu(\mathcal{O})$, up to an explicit degree shift, as graded $R_\alpha(\mathcal{O})$ -modules for any integral domain $\mathcal{O}$.

Recall that $\{v^T \mid T \in \text{St}(\mu)\}$ is a basis of $S^\mu(\mathcal{O})$ and that $\{v_T \mid T \in \text{St}(\mu)\}$ is a basis of $S_\mu(\mathcal{O})$. Let $\{f_T \mid T \in \text{St}(\mu)\}$ and $\{f^T \mid T \in \text{St}(\mu)\}$ be the corresponding dual bases of $S^\mu(\mathcal{O})^\circledast$ and $S_\mu(\mathcal{O})^\circledast$, respectively, so that

$$f_S(v^T) = \delta_{S,T} = f^S(v_T),$$

where $S, T \in \text{St}(\mu)$. By definition, $\deg f_T = -\deg v^T = -\deg T$ and $\deg f^T = -\text{codeg } T$. Recalling (2.14), we now have

$$\deg f_T = \text{codeg } T - \text{def } \alpha \quad \text{and} \quad \deg f^T = \deg T - \text{def } \alpha. \quad (7.23)$$

Lemma 7.24. *As R_α -modules, $S^\mu(\mathcal{O})^\circledast$ is generated by $f_{\mathbf{T}_\mu}$ and $S_\mu(\mathcal{O})^\circledast$ is generated by $f^{\mathbf{T}_\mu}$.*

Proof. We only prove that $S^\mu(\mathcal{O})^\circledast = R_\alpha f_{\mathbf{T}_\mu}$. The proof of the second statement is similar.

We claim that if $T \in \text{St}(\mu)$ then there exist scalars $c_S \in \mathcal{O}$ such that

$$f_T = \psi_T f_{\mathbf{T}_\mu} + \sum_{S \in \text{St}(\mu)} c_S f_S,$$

where $c_S \neq 0$ only if $\ell(w^S) > \ell(w^T)$. The claim implies that $f_T \in R_\alpha f_{\mathbf{T}_\mu}$, for all $T \in \text{St}(\mu)$, so that $S^\mu(\mathcal{O})^\circledast = R_\alpha f_{\mathbf{T}_\mu}$ by the remarks above.

To prove the claim we argue by downwards induction on the dominance order. If $T = \mathbf{T}_\mu$ then $w_{\mathbf{T}_\mu} = 1 = \psi_{\mathbf{T}_\mu}$ so that indeed $f_{\mathbf{T}_\mu} = \psi_{\mathbf{T}_\mu} f_{\mathbf{T}_\mu}$. Next suppose that $T \triangleleft \mathbf{T}_\mu$ and let $S \in \text{St}(\mu)$. Then, by definition,

$$(\psi_T f_{\mathbf{T}_\mu})(v^S) = f_{\mathbf{T}_\mu}(\psi_{w_T^{-1}} v^S) = f_{\mathbf{T}_\mu}(\psi_{w_T^{-1}} \psi_{w^S} z^\mu).$$

By Lemma 2.18(iii), $w^{\mathbf{T}_\mu} = w_T^{-1} w^T$ and $\ell(w^{\mathbf{T}_\mu}) = \ell(w_T^{-1}) + \ell(w^T)$. Consequently, if $\ell(w^S) \leq \ell(w^T)$ and $S \neq T$ then $w^{\mathbf{T}_\mu}$ can not appear as a subexpression of $w_T^{-1} w^S$ so that $(\psi_T f_{\mathbf{T}_\mu})(v^S) = 0$ by Proposition 3.15. Therefore,

the coefficient of f_S in $\psi_T f_{T_\mu}$ is zero whenever $\ell(w^S) < \ell(w^T)$. Finally, consider the case when $S = T$. By Proposition 3.15, there exist polynomials $p_u(y) \in \mathcal{O}[y_1, \dots, y_d]$ such that

$$\begin{aligned} \psi_{w_T^{-1}} \psi_{w^T} z^\mu &= \psi^{T^\mu} z^\mu + \sum_{u < w^{T^\mu}} \psi_u p_u(y) z^\mu \\ &= v^{T^\mu} + \sum_{u < w^{T^\mu}} p_u \psi_u z^\mu, \end{aligned}$$

where $p_u = p_u(0) \in \mathcal{O}$ and the last equality follows from Lemma 6.19. It follows that $(\psi_T f_{T_\mu})(v^T) = 1$. Hence, if we write $\psi_T f_{T_\mu}$ with respect to the basis $\{f_S\}$ then f_T appears with coefficient 1. This completes the proof of the claim and, hence, of the lemma. $\square$

We can now prove the main result of this section.

Theorem 7.25. *As graded $R_\alpha(\mathcal{O})$ -modules,*

$$S^\mu(\mathcal{O}) \cong S_\mu(\mathcal{O})^{\otimes} \langle \text{def } \alpha \rangle \quad \text{and} \quad S_\mu(\mathcal{O}) \cong S^\mu(\mathcal{O})^{\otimes} \langle \text{def } \alpha \rangle$$

Proof. The two isomorphisms are equivalent so we consider only the first isomorphism. By Lemma 7.24 and (7.23) it is enough to show that f^{T^μ} satisfies the defining relations from Definition 5.9 for the element z^μ as, taking into account our basis results, this will imply that there is a unique isomorphism $S^\mu(\mathcal{O}) \xrightarrow{\sim} S_\mu(\mathcal{O})^{\otimes} \langle \text{def } \alpha \rangle$ which sends z^μ to f^{T^μ} . From the definitions, $e(i) f^{T^\mu} = \delta_{ii^\mu}$, so it remains to show that

- (ii) $y_r f^{T^\mu} = 0$ for all $r = 1, \dots, d$;
- (iii) $\psi_r f^{T^\mu} = 0$ for all $r = 1, \dots, d-1$ such that $r \rightarrow_{T^\mu} r+1$;
- (iv) $g^A f^{T^\mu} = 0$ for all row Garnir nodes A in μ .

By freeness it is sufficient to consider the case when $\mathcal{O} = \mathbb{Z}$ and, since $S_\mu(\mathbb{Z})^{\otimes}$ embeds into $S_\mu(\mathbb{C})^{\otimes}$, it is enough to verify the relations when $\mathcal{O} = \mathbb{C}$.

As in Section 6.6, let $\xi = \exp(2\pi i/e)$ if $e > 0$ and if $e = 0$ take ξ to be any non-root of unity in $\mathbb{C}$. Then, by Theorem 6.18, $R_\alpha^\Lambda(\mathbb{C}) \cong H_\alpha^\Lambda(\mathbb{C}, \xi)$, so we can invoke results from [11]. Hence, as graded R_α -modules,

$$\begin{aligned} S^\mu(\mathbb{C}) &\cong S_H^\mu(\mathbb{C}), && \text{by Theorem 6.23,} \\ &\cong S_\mu^H(\mathbb{C})^{\otimes} \langle \text{def } \alpha \rangle, && \text{by [11, Proposition 6.19],} \\ &\cong S_\mu(\mathbb{C})^{\otimes} \langle \text{def } \alpha \rangle, && \text{by Theorem 7.19.} \end{aligned}$$

To complete the proof we scrutinize the second isomorphism above.

In our notation, the proof of [11, Proposition 6.19] shows that there exists a homogeneous associative bilinear form

$$\{ , \} : S_H^\mu \times S_\mu^H \langle \text{def } \alpha \rangle \longrightarrow \mathbb{C}; \quad (a, b) \mapsto \{a, b\},$$

such that $\{v_H^S, v_T^H\} = 0$ unless $T \supseteq S$. (When comparing our notation with [11] the reader should remember that $S_\mu^\mu = S_{\mu'}^{H^M}$ as defined in section 7.8.)

The isomorphism $S_H^\mu(\mathbb{C}) \xrightarrow{\sim} S_\mu^H(\mathbb{C})^{\otimes} \langle \text{def } \alpha \rangle$ is then the map which sends $a \in S_H^\mu(\mathbb{C})$ to $\varphi_a \in S_\mu^H(\mathbb{C})^{\otimes} \langle \text{def } \alpha \rangle$, where $\varphi_a(b) = \{a, b\}$, for all $b \in S_\mu^H(\mathbb{C})^{\otimes} \langle \text{def } \alpha \rangle$. Observe that the triangularity of the form $\{ , \}$ implies that $\varphi_{v^{T^\mu}}$ is a scalar multiple of f^{T^μ} . Therefore, since the map $a \mapsto \varphi_a$ is an isomorphism, it follows from Definition 5.9 and Corollary 7.22 that $\varphi_{v^{T^\mu}}$,

and hence $f^{\mathbf{T}^\mu} \in S_\mu(\mathbb{C})^\otimes$, satisfies the three relations (ii)–(iv) above. Consequently, $f^{\mathbf{T}^\mu} \in S_\mu(\mathbb{Z})^\otimes$ also satisfies these relations and the theorem is proved. $\square$

Remark 7.26. In principle, it should be possible to prove Theorem 7.25 directly by verifying that $f^{\mathbf{T}^\mu}$ satisfies the relations in Definition 5.9. This appears to be an involved calculation.

8. TWO APPLICATIONS

In this section we work again over an arbitrary commutative unital ring $\mathcal{O}$.

8.1. Specht modules for higher levels as induced modules. Let $\mu = (\mu^{(1)}, \dots, \mu^{(l)}) \in \mathcal{P}_\alpha^\kappa$ with $\alpha^{(m)} = \text{cont}(\mu^{(m)})$, for $m = 1, \dots, l$. Then $\alpha = \alpha^{(1)} + \dots + \alpha^{(l)}$. Consider each partition $\mu^{(m)}$ as an element of $\mathcal{P}_{\alpha^{(m)}}^{(k_m)}$; that is, as a partition whose $(1,1)$ -node has residue k_m . Then we have the universal graded Specht modules $S^{\mu^{(m)}}$ for the algebras $R_{\alpha^{(m)}}^{\Lambda_{k_m}}$, for $m = 1, \dots, l$. Inflating along the surjection $R_{\alpha^{(m)}} \twoheadrightarrow R_{\alpha^{(m)}}^{\Lambda_{k_m}}$ we may consider $S^{\mu^{(m)}}$ as a graded $R_{\alpha^{(m)}}$ -module. Note that this graded module is generated by the element $z^{\mu^{(m)}}$ of degree $\deg(\mathbf{T}^{\mu^{(m)}})$.

As in (3.19), consider the graded R_α -module $S(\mu^{(1)}) \circ \dots \circ S(\mu^{(l)})$ which is generated by the element

$$z^{\mu^{(1)}, \dots, \mu^{(l)}} := 1 \otimes (z^{\mu^{(1)}} \otimes \dots \otimes z^{\mu^{(l)}}) \quad (8.1)$$

of degree $\deg \mathbf{T}^{\mu^{(1)}} + \dots + \deg \mathbf{T}^{\mu^{(l)}}$.

Our new definition of Specht modules by generators and relations makes the following useful result almost obvious. Note that in [28], [5, (3.24)] the right hand side of (8.3) was taken as the definition of the Specht module.

Theorem 8.2. *Suppose that $\mu = (\mu^{(1)}, \dots, \mu^{(l)}) \in \mathcal{P}_\alpha^\kappa$. Then*

$$S^\mu \cong S^{\mu^{(1)}} \circ \dots \circ S^{\mu^{(l)}} \langle d_\mu \rangle, \quad (8.3)$$

where

$$d_\mu := \deg(\mathbf{T}^\mu) - \deg(\mathbf{T}^{\mu^{(1)}}) - \dots - \deg(\mathbf{T}^{\mu^{(l)}}).$$

as graded R_α -modules. In particular, $S^{\mu^{(1)}} \circ \dots \circ S^{\mu^{(l)}} \langle d_\mu \rangle$ factors through the surjection $R_\alpha \twoheadrightarrow R_\alpha^\Lambda$, and the isomorphism (8.3) is also an isomorphism of graded R_α^Λ -modules.

Proof. The vector $z^{\mu^{(1)}, \dots, \mu^{(l)}}$ from (8.1) satisfies the defining relations on the vector $z^\mu \in S^\mu$ from Definition 5.9. This yields a homogeneous module homomorphism $S^\mu \rightarrow S^{\mu^{(1)}} \circ \dots \circ S^{\mu^{(l)}} \langle d_\mu \rangle$ which maps z^μ onto $z^{\mu^{(1)}, \dots, \mu^{(l)}}$. To construct the inverse homomorphism, by Frobenius Reciprocity, it suffices to construct a homomorphism of $R_{\alpha^{(1)}, \dots, \alpha^{(l)}}$ -modules

$$S^{\mu^{(1)}} \boxtimes \dots \boxtimes S^{\mu^{(l)}} \rightarrow \text{Res}_{\alpha^{(1)}, \dots, \alpha^{(l)}}^\alpha S^\mu,$$

which maps $z^{\mu^{(1)}} \otimes \dots \otimes z^{\mu^{(l)}}$ onto z^μ . Such homomorphism arises by Definition 5.9 again, using defining relations for the modules $S^{\mu^{(1)}}, \dots, S^{\mu^{(l)}}$. $\square$

8.2. Column Specht modules as signed row Specht modules. In this final section we investigate the analogue of tensoring the Specht modules with the sign representation. Recall the isomorphism $\mathbf{sgn} : R_\alpha \rightarrow R_{\alpha'}$ from (3.14) and the $\mathbf{sgn}$ -twist $M^{\mathbf{sgn}} \in R_\alpha\text{-Mod}$ of a module $M \in R_{\alpha'}\text{-Mod}$. We determine what happens to the Specht modules of $R_{\alpha'}$ under this twist.

For each $\mu \in \mathcal{P}_\alpha^\kappa$ we have row Specht module S^μ and column Specht module S_μ with bases $\{v^{\mathbf{T}}\}$ and $\{v_{\mathbf{T}}\}$, respectively, parametrized by $\mathbf{T} \in \text{St}(\mu)$. Similarly, for each $\nu \in \mathcal{P}_{\alpha'}^{\kappa'}$ we have row Specht module S^ν and column Specht module S_ν , with bases $\{v^{\mathbf{S}}\}$ and $\{v_{\mathbf{S}}\}$, respectively, parametrized by $\mathbf{S} \in \text{St}(\nu)$. The definition of these modules and bases depends on κ and κ' , respectively.

In section 2.2 we defined the conjugate multipartition $\mu' \in \mathcal{P}_{\alpha'}^{\kappa'}$ of the multipartition $\mu \in \mathcal{P}_\alpha^\kappa$. Recall from section 2.3 that the definition of degree and codegree of a tableau $\mathbf{T}$ depends on κ . We write $\deg^\kappa(\mathbf{T})$, $\text{codeg}^\kappa(\mathbf{T})$, $\text{res}^\kappa A$, etc., when we want to emphasize dependence on κ . Finally, the conjugate tableau $\mathbf{T}'$ is defined in section 2.3, and if $\mathbf{i} \in I^\alpha$ then $-\mathbf{i} \in I^{\alpha'}$ is defined in (3.13).

For any node $A = (a, b, m)$ define $A' := (b, a, l - m + 1)$. Note that $A \in \mu$ if and only if $A' \in \mu'$, in which case $\mathbf{T}(A) = \mathbf{T}'(A')$. Moreover, we have $\text{res}^\kappa A \equiv -\text{res}^{\kappa'} A'$ by (2.6), and A is above B if and only if A' is below B' . The following lemma now follows from definitions.

Lemma 8.4. *Suppose that $\mu \in \mathcal{P}_\alpha^\kappa$ and $\mathbf{T} \in \text{St}(\mu)$. Then $\mathbf{i}^{\kappa'}(\mathbf{T}') = -\mathbf{i}^\kappa(\mathbf{T})$, $\deg^\kappa(\mathbf{T}) = \text{codeg}^{\kappa'}(\mathbf{T}')$ and $\text{codeg}^\kappa(\mathbf{T}) = \deg^{\kappa'}(\mathbf{T}')$.*

The main result of this section is:

Theorem 8.5. *Suppose that $\alpha \in Q_+$ with $d = \text{ht}(\alpha)$, and $\mu \in \mathcal{P}_\alpha^\kappa$. Then*

$$S^\mu \cong (S_{\mu'})^{\mathbf{sgn}} \quad \text{and} \quad S_\mu \cong (S^{\mu'})^{\mathbf{sgn}}$$

as graded $R_\alpha(\mathcal{O})$ -modules.

Proof. We claim that there are degree zero homomorphisms of graded $R_\alpha(\mathcal{O})$ -modules

$$\theta^\mu : S^\mu \longrightarrow (S_{\mu'})^{\mathbf{sgn}} \quad \text{and} \quad \theta_{\mu'} : (S_{\mu'})^{\mathbf{sgn}} \longrightarrow S^\mu$$

such that $\theta^\mu(z^\mu) = z_{\mu'}$ and $\theta_{\mu'}(z_{\mu'}) = z^\mu$. As z^μ and $z_{\mu'}$ generate the two Specht modules, this claim implies the theorem.

Note that

$$\deg z^\mu = \deg^\kappa(\mathbf{T}^\mu) = \text{codeg}^{\kappa'}(\mathbf{T}_{\mu'}) = \deg z_{\mu'}$$

by Lemma 8.4. So to prove the existence of θ^μ , it suffices to check that $z_{\mu'} \in (S_{\mu'})^{\mathbf{sgn}}$ satisfies the defining relations of S^μ from Definition 5.9. The map $\theta_{\mu'}$ is constructed similarly using Definition 7.11 instead, so we only give details for θ^μ .

By Lemma 8.4, $\mathbf{i}^\mu = \mathbf{i}(\mathbf{T}^\mu) = -\mathbf{i}(\mathbf{T}_{\mu'}) = -\mathbf{i}_{\mu'}$. Therefore, if $\mathbf{j} \in I^\alpha$ then

$$e(\mathbf{j}) \cdot z_{\mu'} = e(-\mathbf{j})z_{\mu'} = \delta_{-\mathbf{j}, \mathbf{i}_{\mu'}} z_{\mu'} = \delta_{-\mathbf{j}, -\mathbf{i}^\mu} z_{\mu'} = \delta_{\mathbf{j}, \mathbf{i}^\mu} z_{\mu'}.$$

Therefore, $z_{\mu'}$ satisfies Definition 5.9(i). Moreover, $y_r \cdot z_{\mu'} = -y_r z_{\mu'} = 0$ for all $1 \leq r \leq d$. Next observe that if $1 \leq r < d$ then $r \rightarrow_{\mathbf{T}^\mu} r+1$ if and only if $r \downarrow_{\mathbf{T}_{\mu'}} r+1$. Hence, $\psi_r \cdot z_{\mu'} = -\psi_r z_{\mu'} = 0$, by Definition 5.9(iii) and Definition 7.11(iii).

It remains to check that $z_{\mu'} \in (S_{\mu'})^{\text{sgn}}$ satisfies the row Garnir relations from Definition 5.9(iv). Recall the node correspondence $A \leftrightarrow A'$ defined before Lemma 8.4 which sends a node $A \in \mu$ to $A' \in \mu'$. If $A \in \mu$ is a row Garnir node then $A' \in \mu'$ is a column Garnir node and, further, this correspondence sends row bricks in μ to column bricks in μ' . In particular, $k^A = k_{A'}$, where k^A is the number of row bricks in $\mathbf{B}^A$ and $k_{A'}$ is the number of column bricks in $\mathbf{B}_{A'}$. Moreover, $\text{sgn}(\tau_r^A) = \tau_{A'}^r$ by Lemma 7.3(i), for $1 \leq r < k_A$, so that $\text{sgn}(g^A) = g_{A'}$. Therefore,

$$g^A \cdot z_{\mu'} = \text{sgn}(g^A) z_{\mu'} = g_{A'} z_{\mu'} = 0,$$

where the last equality is a column Garnir relation in $S_{\mu'}$. $\square$

REFERENCES

- [1] S. Ariki, On the decomposition numbers of the Hecke algebra of $G(m, 1, n)$, *J. Math. Kyoto Univ.* **36** (1996), 789–808.
- [2] J. Brundan, Centers of degenerate cyclotomic Hecke algebras and parabolic category $\mathcal{O}$, *Represent. Theory* **12** (2008), 236–259.
- [3] J. Brundan and A. Kleshchev, Blocks of cyclotomic Hecke algebras and Khovanov-Lauda algebras, *Invent. Math.* **178** (2009), 451–484.
- [4] J. Brundan and A. Kleshchev, Graded decomposition numbers for cyclotomic Hecke algebras, *Adv. Math.* **222** (2009), 1883–1942.
- [5] J. Brundan and A. Kleshchev, The degenerate analogue of Ariki’s categorification theorem, *Math. Z.* **266** (2010), 877–919.
- [6] J. Brundan, A. Kleshchev and W. Wang, Graded Specht modules, [arXiv:0901.0218](https://arxiv.org/abs/0901.0218), to appear in *J. reine angew. Math.*
- [7] R. Dipper, G. D. James and A. Mathas, Cyclotomic q -Schur algebras, *Math. Z.* **229** (1998), 385–416.
- [8] H. Garnir, Théorie de la représentation lineaire des groupes symétriques, *Mém. Soc. Roy. Sci. Liège (4)* **10** (1950).
- [9] J. J. Graham and G. I. Lehrer, Cellular algebras, *Invent. Math.* **123** (1996), 1–34.
- [10] I. Grojnowski, Affine $\mathfrak{sl}_p$ controls the representation theory of the symmetric group and related Hecke algebras, [arXiv:math.RT/9907129](https://arxiv.org/abs/math.RT/9907129).
- [11] J. Hu and A. Mathas, Graded cellular bases for the cyclotomic Khovanov-Lauda-Rouquier algebras of type A, *Adv. Math.*, **225** (2010), 598–642.
- [12] J. Hu and A. Mathas, Graded induction for Specht modules; [arXiv:1008.1462](https://arxiv.org/abs/1008.1462).
- [13] G. D. James, *The representation theory of the symmetric groups*, Lecture Notes in Mathematics 682, Springer, Berlin, 1978.
- [14] G. D. James and A. Mathas, The Jantzen sum formula for cyclotomic q -Schur algebras, *Trans. Amer. Math. Soc.* **352** (2000), 5381–5404.
- [15] V. G. Kac, *Infinite Dimensional Lie Algebras*, Cambridge University Press, 1990.
- [16] M. Khovanov and A. Lauda, A diagrammatic approach to categorification of quantum groups I, *Represent. Theory* **13** (2009), 309–347.
- [17] M. Khovanov and A. Lauda, A diagrammatic approach to categorification of quantum groups II; [arXiv:0804.2080](https://arxiv.org/abs/0804.2080).
- [18] A. Kleshchev, *Linear and Projective Representations of Symmetric Groups*, Cambridge University Press, Cambridge, 2005.
- [19] A. Kleshchev, Representation theory of symmetric groups and related Hecke algebras, *Bull. Amer. Math. Soc.* **47** (2010), 419–481.
- [20] S. Lyle and A. Mathas, Blocks of cyclotomic Hecke algebras, *Advances Math.* **216** (2007), 854–878.
- [21] A. Mathas, *Iwahori-Hecke algebras and Schur algebras of the symmetric group*, University Lecture Series 15, American Mathematical Society, Providence, RI, 1999.
- [22] A. Mathas, Matrix units and generic degrees for the Ariki-Koike algebras, *J. Algebra*, **281** (2004), 695–730.

- [23] G. E. Murphy, On the representation theory of the symmetric groups and associated Hecke algebras, *J. Algebra* **152** (1992), 492–513.
- [24] A. Ram, Skew shape representations are irreducible, 161–189 in *Combinatorial and geometric representation theory (Seoul, 2001)*, Contemp. Math., **325**, Amer. Math. Soc., Providence, RI, 2003.
- [25] R. Rouquier, 2-Kac-Moody algebras; [arXiv:0812.5023](https://arxiv.org/abs/0812.5023).
- [26] W. Specht, Die irreduziblen Darstellungen der symmetrischen Gruppe, *Math. Z.* **39** (1935), 696–711.
- [27] J.R. Stembridge, On the fully commutative elements of Coxeter groups, *J. Algebraic Combin.* **5** (1996), 353–385.
- [28] M. Vazirani, Parameterizing Hecke algebra modules: Bernstein-Zelevinsky multisegments, Kleshchev multipartitions, and crystal graphs, *Transform. Groups* **7** (2002), 267–303.
- [29] A. Young, On the quantitative substitutional analysis I, *Proc. London Math. Soc.* (1) **33** (1900), 97–145.
- [30] A. Young, On the quantitative substitutional analysis II, *Proc. London Math. Soc.* (1) **34** (1901), 361–397.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OR 97403, USA
E-mail address: `klesh@uoregon.edu`

SCHOOL OF MATHEMATICS AND STATISTICS F07, UNIVERSITY OF SYDNEY, NSW 2006, AUSTRALIA
E-mail address: `andrew.mathas@sydney.edu.au`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MELBOURNE, VIC 3010, AUSTRALIA
E-mail address: `aram@unimelb.edu.au`